KINGDOM OF SAUDI ARABIA
**Northern Border University (NBU)**
**Journal of the North for Basic & Applied Sciences( JNBAS)**
**p - ISSN :** 1658 -7022 / **e-ISSN:** 1658 - 7014
www.nbu.edu.sa
s.journal@nbu.edu.sa

# Phytobezoar Predisposing Factors: Re-evaluating the Role of Dietary Fibres - A Systematic Review

**Mohamed Ashour**
**Department of pharmacognosy and alternative medicine, faculty of Pharmacy, Northern Border University, Rafha, 91911 Saudi Arabia**

**Ahmed Ashour**
**Department of cardiothoracic surgery, faculty of medicine, Ain shams University, Cairo, Egypt.**

(**Received:** :3rd May 2026; **Accepted:** 14th June 2026)

**Abstract**
**Background:** Phytobezoar is defined as the compaction of indigestible food material (mainly fibrous plant tissues) localised in the gastrointestinal tract (GIT), leading mainly to GI obstruction (GIO) and many GI health problems. Some researchers claimed that vegetables, fruits, and other DF-containing foods are the main culprits in phytobezoar formation.
**Objective:** systematic review of the available case reports and case series to identify the actual culprits of phytobezoar formation and to distinguish the role of dietary fibres.
**Data Source:** We searched PubMed, MEDLINE, Web of Science platform, and Google Scholar with no restriction through April 2026.
**Study Selection:** available phytobezoar cases (including reports, series and reviews).
**Data Extraction:** We extracted data on patient demographic information, diagnosis, and emphasizing the predisposing/risk factors.
**Data Synthesis:** 218 publications met inclusion criteria, reporting on 1382 cases. The cases have been categorized according to the predisposing/ risk factors.
**Limitations:** no limitation for time, age, and geography, however the study is limited to English publications.
**Conclusion:** the study unambiguously distinguished the actual predisposing/risk factors for phytobezoar formation including previous GI surgeries, unusual foodstuffs, among others. Although phytobezoars are usually formed of undigested fibrous plant material, however, proper consumption of the recommended amounts of dietary fibre-containing foods is not responsible for phytobezoar formation.

**Keywords:** Dietary fibres, phytobezoar, bezoar culprits, gastrointestinal surgery.

1658-7022© JNBAS. (1448 H/2026). Published by Northern Border University (NBU). All Rights Reserved.

(*) **Corresponding Author**:
Mohamed Ashour
Department of pharmacognosy and alternative medicine, faculty of Pharmacy, Northern Border University, Rafha, 91911 Saudi Arabia

**E-mail:**Ashourmohamed08@yahoo.com
**ORCID ID:** 0000-0002-1298-5885
**Permanent Address:** Department of Pharmacognosy, Faculty of Pharmacy, Al-Azhar University, Nasr City 11884, Cairo, Egypt.

**DOI:** 10.12816/0062625

## 1. Introduction

The term "Phytobezoar" has been derived from the Greek word "phyton" (plant) and "bezoar" is thought to refer to the Persian words *pâd* (expelling) and *zahr* (poison).[1,2] The medical term "bezoar" was established in the medical literature since 1854 by Dr. Richard Quain.[3] In medical terminology, bezoars represent an agglomerate of foreign materials, hairs (trichobezoars), indigestible plant material (phytobezoars), or drugs (pharmacobezoars) that ingested, accumulate and solidify inside GIT, creating a solid phase that progressively grows until the point it can prevent the normal digestive process and often obstructs GIT.[3-6]

Although rare cause of primary gastrointestinal obstruction, phytobezoars are the most common cause of bezoar occurrence, accounting for up to 70% of all bezoars.[4,7,8]

Bezoars are responsible for 0.4% - 4% of cases of mechanical gastrointestinal obstruction worldwide. [9,10] Some researchers have claimed that vegetables, fruits, and other fibre-containing foods are the main culprits in the formation of phytobezoars.[3,5,6,11,12] However, they do not differentiate between the recommended amount of dietary fibres and other fibrous plant tissues (unusual food material) that are consumed intentionally or accidentally, leading to phytobezoar formation. Dietary fibres (DF) is a term used to describe indigestible, soluble and insoluble part of usual foodstuffs, especially carbohydrates.[13] These DF are found naturally at a reasonable level in our healthy foods e.g. popular fruits and vegetables, either as soluble fibres (e.g. mucilaginous materials: psyllium seed, chia seeds, Okra, fenugreek, salep tubers, grains …etc.) or insoluble fibres (e.g. grain walls (wheat Bran, oat bran), Beans, shelled nuts, vegetables, fruits ...etc.).[14-17] Here, we have to recognise non-food plant fibres e.g. plant wood fibers, hard or leathery seed skins (sunflower seed coats, watermelon seed coat, Apricot pits, etc.,), these types of plant fibres are not considered as DF. According to common recommendations, DF should be consumed regularly as food habit at a daily dose of 30-38 g for adult men as a part of 2600 kcal food/day, and 28-30 g for adult women as part of 2000 kcal daily food.[13-16] Dieticians and healthcare providers must consider these values in their prescriptions and recommendations to the public. They must also recalculate these quantities based on patient's age and health status. Consumption of the recommended amount of DF has a positive impact and offers great health benefits. DF consumption was linked to increased insulin sensitivity,[18] improved blood sugar levels,[18] management of diabetes,[19]improvement of obesity and body mass index (BMI),[18] prevention of constipation[20], lower blood cholesterol levels,[21] decreased fat and carbohydrate absorption,[22]and lowering of blood pressure.[23,24] Other benefits have also been confirmed, such as improvement of GIT movement ,[22] and increased mineral absorption in the intestinal tract,[25] reduces energy intake,[26] decreases the chance of pathogenic microbial growth, increases the chance of growth and development of beneficial microbial flora,[27] biosynthesis of short-chain fatty acids with valuable health benefits,[28] improvement of the immune system,[29] decreases the absorption of toxic substances and carcinogens for GIT,[30] and prevention of tumour formation (e.g. colorectal cancer) and consequently, enhances the quality of life.[13,31,32]

### 1.1 Mechanisms of phytobezoar formation:

Phytobezoars are formed in the alimentary canal as a result of specific indigestible plant tissues and biological reactions that interact to form solid or semisolid concretions or conglomerates that are difficult to hydrolyze or fragment and cause partial or complete GIO. Two scenarios have been accepted as mechanisms for phytobezoar formation.

- **Chemical scenario:** Phytobezoar formation (diospyrobezoars) due to consumption of persimmon fruits (*Diospyros kaki*) has been confirmed. Unripe persimmon fruits contain high concentrations of tannins (phlobatannin: shibuol), fibre contents, and indigestible soluble carbohydrates.[33,34] When tannin comes into contact with gastric juice (higher gastric acidity) in the stomach, especially in an empty stomach, it can precipitate and undergo polymerization, forming a coagulum that includes cellulose, hemicellulose, and protein. This coagulum serves as the foundation for the bezoar. As this cluster continues to evolve, it undergoes dehydration and solidification processes. This solidified mass grows, resists fragmentation or hydrolysis and can then be moved to narrow spaces in the GIT, leading to obstruction.[7,8,34]
- **Mechanical scenario:** the formation of phytobezoar may be due to accumulation of indigestible plant materials such as hard leathery or woody plant material (e.g. unshelled watermelon seeds, unshelled pumpkin seeds), or a high amount of hygroscopic material (e.g. dry chia seeds), where these materials accumulate and grow to form a nucleus of phytobezoar under the influence of predisposing factors such as GIT strictures or GI laziness.[35] However, other scenarios have not been confirmed for phytobezoar formation due to easily digestible food materials such as rice, potato-based foods, and meat boluses.[36]

### 1.2 Phytobezoar - related complications and health problems:

Gastrointestinal obstruction (GIO) is the main complication of phytobezoar formation, where the

phytobezoar can be localised in the stomach and move to close the pyloric canal or continue moving to cause small bowel obstruction in the duodenum, jejunum, or ileum, preventing the normal function of the GIT.[3] However, seed shell-related phytobezoars are commonly formed or agglomerated in the large intestine.[37-39] Gastrointestinal (GI) obstruction of the GIT can lead to serious health issues, such as GI perforation, GI bleeding, intussusception, and strangulation, which may lead to local ischaemia and necrosis.[6,40] Indeed, acute pancreatitis[41], Jejunal Obstruction[42], GI Hemorrhage[43], Restrictive lung pattern[44], small bowel perforation[45], Proximal Ileal Perforation[46], body weight loss, GI ulceration, GI inflammation, fistula formation, and adhesion have also been reported as complications for phytobezoars.[47]

phytobezoar cases have been reported in almost all geographic regions, worldwide due to fibrous and less likely non-fibrous plant materials and non-plant foodstuffs. The exact role of dietary fibres in the formation of phytobezoar is debatable. Consequently, an urgent literature review is needed to clarify the actual causes of phytobezoars. The present study was designed to investigate the published phytobezoar cases from global healthcare centres in order to explain the actual culprits of phytobezoar formation, and distinguishes the role of dietary fibres.

## 2. Materials and Methods

### 2.1 The study selection and literature search strategy

The data of the present systematic review were obtained from Life-science databases (PubMed, MEDLINE, Web of Science platform, and Google Scholar).48-51 The search was conducted independently by two reviewers (MA and AA) using specific terms "phytobezoar", "phytobezoar-predisposing factors", "phytobezoar-dietary fibres", and "dietary fibres" to identify studies relevant to phytobezoar cases and the role of dietary fibres. The obtained search results were filtered first by screening titles and abstracts then further filtering has been made to exclude duplicate and irrelevant publications. Then further filtering was performed to exclude irrelevant publications and those lacking important information. Disagreements were resolved by consensus. The search included studies published up to 2026.

### 2.2 Quality assessment of the eligible publications

The assessment was carried out independently by two investigators (MA & AA) addressing the items available in the case reports. These items include 1) relevance of the title to the reported information, 2) adequacy of demographic information, 3) relevant past history, 4) case diagnosis, 5) type of concretion (phytobezoar mass), and 6) pre-disposing/risk factors if any, 7) discussion of the relevant evidences. Possible item ratings are yes, partially, or no; disagreement was resolved by consensus.

### 2.3 Inclusion and Exclusion criteria

**Inclusion criteria:** All studies reported human cases of phytobezoar. There was no limitation for time of publication, and no limitation of countries or geographic regions.

**Exclusion criteria:** 1) non-plant-related bezoar cases (e.g. trichobezoars, pharmacobezoars); 2) all articles that are free from reported phytobezoar information; 3) non-human phytobezoar cases (e.g. animal bezoar cases); 4) repetitions or case duplicates were also excluded; 5) lacking of important information. 6) articles published in languages other than English.

### 2.4 Data synthesis and analysis

218 articles have been selected based on the eligibility criteria for inclusion and exclusion, Descriptive statistical analyses were used to data syntheses and categorization process. Microsoft Excel software (Microsoft, USA) has been utilized for Statistical analysis, classification, designing graphs and extracting tables.

## 3. Results

### 3.1 Quality appraisal

Based on the items of quality assessment, the overall quality of the included studies was moderate to high. The titles were relevant to the reported information (100%), adequacy of demographic information (age, gender, nationality, residency) (80.4%), relevant past history (93%), case diagnosis (99.0%), type of concretion (phytobezoar mass) (67.9%), pre-disposing/risk factors if any (89.9%), and discussion of the relevant evidences (98.8%).

### 3.2 Publication characteristics

The eligible publications (218 articles), have been investigated (Figure 1). These eligible publications were found to comprise almost all relevant topics, almost all geographic regions (worldwide), with no limited time period through 2026. The selected articles included case reports, case series, case reports with reviews, review of cases, and other relevant reviews of phytobezoar occurrence. The eligible records were carefully reviewed, and extracted to the Microsoft excel sheets, then categorized into groups as seen in the provided tables (S1 appendix). Total Number of reviewed phytobezoar cases (1382 cases) have been categorised according to patient age with or without previous GI surgery (Figure 2). The phytobezoar cases without previous GI surgery have been classified according to predisposing/risk factors (Figures 3).





**Figure 1. PRISMA 2020 Flow diagram for the selection of records and publications.**





**Figure 2. Classification of phytobezoar cases according to age and previous GI surgery.**





**Figure 3. Classification of phytobezoar cases not associated with previous GI surgery.**

### 3.3 Phytobezoar predisposing factors

Several predisposing/risk factors have been recognised (see supporting information tables 1 -6).[34,35] The most common predisposing/risk factors (Figure 4) have been considered the main culprits and discussed in detail in the following sections.





**Figure 4: The main culprits for phytobezoar formation**

### 3.4 The Actual culprits in phytobezoar formation

#### 3.4.1 Previous GI surgery

Regardless of the nature of the foodstuff, previous GI surgery represents the major risk factor for phytobezoar formation.[52-59] The present study confirmed the presence of 457 phytobezoar cases (33.06 % of all reviewed cases) of different food materials associated with previous GI surgery (S1 appendix - table 1).

### 3.4.2 Unripe Persimmon ingestion

In the present study, we have found 296 (diospyrobezoars) cases (21.41 % of total reported cases) due to persimmon ingestion,[8,35,60,61,62] of which 88 cases associated with previous GI surgery while 208 cases not associated with previous surgery, (S1 appendix - table 2). However, Izumi et al. suggested a widely accepted mechanism (chemical scenario, mentioned above) for diospyrobezoar formation based on gastric acidity and the chemical nature of unripe persimmon fruits which contain a large quantity of tannin (shibuol) to form a sticky coagulum easily converted to a semisolid ball representing the basic nucleus of diospyrobezoar.[34] Thus, diospyrobezoar formation depends on two factors: 1) excessive consumption of unripe persimmon fruits, and 2) concentrated gastric acidity. According to Izumi's hypothesis, the fibre content is not responsible for diospyrobezoar formation. Nonetheless, the United States Department of Agriculture (USDA) report[63] states that the total fibre content of persimmon fruit is only 3.6 g/ 100 g fruits, this equal to only 10% of the recommended daily amount of DF.

### 3.4.3 Unshelled/woody-skin Seeds

44 cases of sunflower seed bezoar, 41 cases of prickly pear bezoar, 66 cases of watermelon seed bezoar, 15 cases of jungle banana seed bezoar, 20 cases of guava seed bezoar, in addition to 49 cases of other seed bezoars including, pumpkin, date, Olive, *Baccaurea macrocarpa*, jaboticaba, sesame, peanuts, lupin, popcorn, pomegranate, tangerine, granadilla, lentil, cherry pits, box myrtle and mango seed bezoars[64-67] have been reported in the present study (S1 appendix - table 3). The unshelled seeds (e.g. sunflower and watermelon seeds) are not usual food; they are ingested with their seed coats (seed skins) which are formed of hard woody or cellulosic shells, and are mainly consumed accidentally or due to psychic disorders.

### 3.4.4 Fruit kernels (Pits)

Hard pits including date and olive pits[38], apricot pits[68,69], nectarine pit[70], prune pit[71], chestnuts[72,73], all have been reported as causes for phytobezoar formation. (S1 appendix - table 3). These types of phytobezoars are due to high cellulosic woody indigestible plant tissues; however, these plant tissues have never been considered dietary fibres and are accidentally ingested forming bezoar and may develop GIO.

### 3.4.5 Dietary fibre-containing herbal medicines without professional instructions

FDA, 2007[74] reported a total of 78 cases of phytobezoar or GIO due to ingestion of laxative drugs formed mainly of psyllium seeds or psyllium seed husks in the period from 1980 to-2000 in the USA.[74] Moreover, Hefny et al., Salguero Molpeceres et al., Agha et al., and shin et al.[75-78] have reported cases of gastrointestinal obstruction due to psyllium seeds and psyllium husks. A similar type of GIO has been reported by Rawl and Brawne due to chia seed ingestion.[79] Similar issues have been reported[80-85] due to the ingestion of Karaya gum, linseed, wheat bran, and oat bran (S1 appendix - table 4). These herbal medicines contain a reasonable amount of dietary fibre (soluble and insoluble). Ingestion of such products must follow the orientation and recommendations of the health authorities, considering both the method of ingestion and the quantity to be ingested. These bulk-forming herbal medicines are strong hygroscopic agents due to their soluble fibre content (mucilages). Psyllium and chia seeds can absorb water equal to 28 times their weights. Therefore, the recommended dose of these herbal drugs must be soaked in sufficient quantity of water for at least 2 Hours prior to ingestion and must be swallowed with plenty of water; otherwise, when ingested dry, they will absorb GI fluid and conglomerate to form a sticky and hard compaction, leading to GIO. Similar cases have been reported due to ingestion of dry psyllium seed husk powder[78], bran[82,84].

### 3.4.6 Poor mastication of Pulpy Fruits and Vegetables

Pulpy fruits and vegetables, such as orange pulp, dry apples, dry apricots, fibrous leaves of rhubarb, and dry grapes[57, 86, 87], have been reported as risk factors for bezoar formation (S1 appendix - table 5). Buchholz and Haisten reported 150 cases of GI obstruction due to different types of phytobezoars (orange pulp, figs, apples, grapefruit, persimmons, coconuts, cabbage, beans, and potatoes) following gastric surgery for duodenal ulcer.[57] They mentioned that 90 % of reported cases were due to orange pulp.[57] They attributed the incidence of phytobezoars to gastric surgery which resulted in loss of normal pyloric function with low gastric acidity, in addition to inadequate chewing of fibrous food, leading to GIO.[57] They published a figure showing approximately intact orange segments as a result of poor mastication. Therefore, inadequate chewing, large amounts of pulpy vegetables and/or fruits, inadequate fluid or water intake and/or previous GI surgery are the main culprits for such bezoar formation.

### 3.4.7 Easily digested food can also produce GIO in certain condition

Many cases of GIO have been reported due to easily-digested foods like rice cakes, mashed potatoes and meat-bolus[36, 57, 88, 89, 90, 91] (S1 appendix - table 5). These reported cases suggested the predisposing factors as aging, improper cooking, quick feeding, and poor mastication as a result of dental disorders.

### 3.4.8 Unusual foods

Katsinelos et al. reported a case of phytobezoar resulting from the ingestion of a substantial quantity of honeycomb in young, healthy Greek woman aged 44 years.[92] Similarly, Hsu et al. documented an unusual

case of phytobezoars due to the consumption of large amounts of vegetable-derived fats (S1 appendix - table 5). It is important to note that these health fads are not containing dietary fibres and are not considered typical food materials.[93]

#### 3.4.9 Dysautonomia and GI laziness

Many phytobezoar cases due to dysautonomia, dysmotility and GI laziness have been reported. (S1 appendix - table 6). These predisposing factors present as a results of GI surgery, GI disorders and/or chronic diseases.[5,7,56,59,94-97]

#### 3.4.10 Psychic disorders

Boopathy and Palaniswamy reported a case of giant bezoar due to accidental ingestion of cotton thread along with different objects such as pieces of bone, seeds of various fruits, stapler pins, buttons, and metal wires for a patient with mental disability.[98]

#### 3.4.11 Aging

Geriatric patients and healthy individuals of old age are more susceptible to phytobezoar formation, regardless of the type of food material. They have many predisposing factors, for example: 1- decreased physical fitness; 2- the expected dental issues and related poor mastication; 3- chronic diseases like diabetes, hypertension, psychic depression, malabsorption, impaired metabolism, in addition to drug-related side effects on GIT.[6] several hundreds of phytobezoar cases have been reported for elderly patients. [6,61, 99-102]

However, phytobezoars can develop in patients without common risk or predisposing factors; therefore, they should not be underestimated, especially when GIO is predicted.[56]

#### 3.4.12 The role of dietary fibres in formation ofPhytobezoar

Many studies have implicated dietary fibres as the main reason or risk factor for phytobezoar formation,[6,13,103] while others have reported a strong association between vegetable and fruit ingestion and phytobezoar formation.[103,104] Although phytobezoars are composed mainly of plant fibres (DF and/or non-DF), However, there are many prominent evidences that confirms the non-liability of the recommended amount of dietary fibres to form phytobezoar unless the predisposing factors are present. The most prominent evidence is the positive response of the majority of people worldwide to the recommendations of Global health authorities to eat a reasonable amount of dietary fibres daily without reporting phytobezoar formation, except very rare cases. These rare cases have been reported for patient having one or more predisposing factors (e.g. previous surgery, GI laziness, chronic diseases, poor mastication, …..etc.).

## 4. Discussion

The present systematic review explained prominently the main culprits that responsible for phytobezoar formation. Although phytobezoar are formed usually of plant fibres (dietary and non-dietary fibres),[5,13,34,56] however, the formation of phytobezoar depends always on the presence of the predisposing factors.[2,4,6,7] Despite of the age, gender, race and geography, phytobezoar formation is expected for any person having the predisposing factors[56]. Prevention of full-beneficial dietary fibres-containing foods is not a proper decision. However, several good practices can be applied to prevent and help manage this health problem. These practices include: a) Health education programs for patients at risk such as those with previous GIT surgery and GIT disorders. Based on various studies, it was established that, maintaining gastrointestinal healthy status through regularisation of meal times and appropriate nutrition, drinking sufficient quantities of water, good mastication, and following health authority recommendations are of great value for preventing bezoar formation. b) parents and relatives of babies and children must be exposed to educational health programs to encourage them to enhance awareness about the predisposing/risk factors. c) herbal medicine practitioners and dieticians must strictly follow and implement the governmental regulations for dispensing herbal medicines and bulk-forming drugs. Indeed, unqualified herbal medicine practitioners must be subjected to an appropriate educational programs. d) the patient and people at risk must be educated about the importance of sport exercise and its positive impact on keeping GI function and prevention of bezoar formation. e) early diagnosis by specialised physicians and effective medical intervention have been considered to avoid or minimise the phytobezoar-related complications.[6, 105, 106,107]

### 4.1 Limitation of the study

- The present study excluded the publications in languages other than English.
- Some records are lacking some information regarding the exact type of phytobezoar.
- Some case reports didn't report the actual predisposing factors, instead they were more interested in diagnosis and management.

## 5. Conclusion

Phytobezoars are annoying health problem, responsible for approximately 4.0 % of GIO, in addition to other GI health issues. Based on the present study, the responsibility of the recommended dietary fibres for phytobezoar formation is not established. the recommended amount and types of dietary fibres have great beneficial effects and cannot be excluded from the healthy foods. However, much attention must be paid for people at risk, and those having one or more predisposing factors.

### 5.1 Data Availability Statement

All data summarized in the text and tables of this review article were obtained from previously published studies with appropriate attributions cited in the text and tables and complete references provided in the reference section.

### 5.2 Supplementary information

- The summary of the reported cases (S1 Appendix)
- PRISMA 2020 checklist (S2 Appendix)

### 5.3 Disclosure statement

- The authors declare no conflicts of interest, and no financial supports.

### 5.4 Author Contributions

- Ashour M: Conceptualisation, research proposal, Data collection, formal analysis, writing-original draft, visualisation; Ashour A: critical discussion, case report review, review-original draft.

### 5.5 Acknowledgement

- The Authors are deeply grateful to Prof. Dr. Hasnaa Abuseif - Dept. of public health and community medicine, Faculty of medicine, Ain Shams university, Egypt - for kind help and support during manuscript revision and validation.

## 6. References

1. *Brightwen, E. (1893). BEZOARS. Science (New York, N.Y.), 21(521), 50–51. https://doi.org/10.1126/science.ns-21.521.50*
2. *Harshberger J. W. (1898). A Review of Our Knowledge of Phyto-Bezoars. The Journal of comparative medicine and veterinary archives, 19(3), 143–145.*
3. *Kosmidis, C. S., Mystakidou, C. M., Varsamis, N., Koulouris, C., Sevva, C., Papadopoulou, K., Michael, C., Katsios, N. I., Theodorou, V., Miltiadous, P., Papadopoulos, K., Vlassopoulos, K., Zarampouka, K., & Mantalovas, S. (2023). Phytobezoar-Induced Mechanical Ileus and Incipient Intussusception: A Case Report. Medicina (Kaunas, Lithuania), 59(7), 1227. https://doi.org/10.3390/medicina59071227*
4. *Gentile, M., Illario, M., De Luca, V., Cestaro, G., Velotti, N., Sivero, S., & Musella, M. (2023). Gastrointestinal bezoars: Review of the literature and report of a rare case of pumpkin seed rectal impaction. Asian journal of surgery, 46(9), 3432–3436. https://doi.org/10.1016/j.asjsur.2023.05.030*
5. *Palchaudhuri, S. (2021). "Bezoars: recognizing and managing these stubborn, sometimes hairy, roadblocks of the gastrointestinal tract." Pract Gastroenterol. Nutrition issues in gastroenterology, series #208 https://med.virginia.edu/ginutrition/wp-content/uploads/sites/199/2021/03/Bezoars-Mrach-2021.pdf*
6. *Yang, S., & Cho, M. J. (2021). Clinical Characteristics and Treatment Outcomes Among Patients With Gastrointestinal Phytobezoars: A Single-Institution Retrospective Cohort Study in Korea. Frontiers in surgery, 8, 691860. https://doi.org/10.3389/fsurg.2021.691860*
7. *Shah, M., Wan, D. Q., Liu, Y., & Rhoads, J. M. (2021). Gastric Bezoar: Retrieve it, Leave it, or Disbelieve it?. Journal of pediatric gastroenterology and nutrition, 72(2), e31–e36. https://doi.org/10.1097/MPG.0000000000002923*
8. *Odeh, A. M., Alkhalifa, A. A., AlHajji, M. A., Alahmed, A. J., Alsalman, J. H., AlMulhem, A. M., AlGhadeer, M. S., Kurdi, E., Albader, F. S., Alsuwaigh, A., & Khan, M. A. (2024). A Rare Case of Gastric Phytobezoar (Diospyrobezoar) in a Healthy Adult. Cureus, 16(8), e68353. https://doi.org/10.7759/cureus.68353*
9. *Dikicier, E., Altintoprak, F., Ozkan, O. V., Yagmurkaya, O., & Uzunoglu, M. Y. (2015). Intestinal obstruction due to phytobezoars: An update. World journal of clinical cases, 3(8), 721–726. https://doi.org/10.12998/wjcc.v3.i8.721*
10. *Guner, A., Kahraman, I., Aktas, A., Kece, C., & Reis, E. (2012). Gastric outlet obstruction due to duodenal bezoar: A case report. International journal of surgery case reports, 3(11), 523–525. https://doi.org/10.1016/j.ijscr.2012.07.005*
11. *Powell, S. D., & Pereira, N. (2023). Bezoars: Culprits of gastrointestinal obstruction that may lead to surgical intervention and potentially surgical complications. Clinical case reports, 11(12), e8126. https://doi.org/10.1002/ccr3.8126*
12. *van Berge Henegouwen, G. P. (1991). "Zijn diëten rijk aan voedingsvezels zinvol?" [Are diets rich in dietary fibre advisable?]. Ned Tijdschr Geneeskd. 135, no. 29: 1306-10.*
13. *Ioniță-Mîndrican, C. B., Ziani, K., Mititelu, M., Oprea, E., Neacșu, S. M., Moroșan, E., Dumitrescu, D. E., Roșca, A. C., Drăgănescu, D., & Negrei, C. (2022). Therapeutic Benefits and Dietary Restrictions of Fiber Intake: A State of the Art Review. Nutrients, 14(13), 2641. https://doi.org/10.3390/nu14132641*
14. *Dragan, S.R., Damian, G., Pah, A.M., Hayta, M., Kahraman, K. (2021). Dietary Fibers. In: Xiao, J., Sarker, S.D., Asakawa, Y. (eds) Handbook of Dietary Phytochemicals. Springer, Singapore. https://doi.org/10.1007/978-981-15-4148-3_31*

15. *Jones J. M. (2013). Dietary fiber future directions: integrating new definitions and findings to inform nutrition research and communication. Advances in nutrition (Bethesda, Md.), 4(1), 8–15. https://doi.org/10.3945/an.112.002907*
16. *McCleary, B. V; Cox, J; Ivory, R; and Delaney, E. (2018). "Definition and Analysis of Dietary Fibre in Grain Products." (CHAPTER 6) In: Cereal Grain-based Functional Foods: Carbohydrate and Phytochemical Components; Beta, T., Camire, M. E. eds.; The Royal Society of Chemistry, (2018); pp. 103-126. https://doi.org/10.1039/9781788012799-00103*
17. *Williams, B. A., Mikkelsen, D., Flanagan, B. M., & Gidley, M. J. (2019). "Dietary fibre": moving beyond the "soluble/insoluble" classification for monogastric nutrition, with an emphasis on humans and pigs. Journal of animal science and biotechnology, 10, 45. https://doi.org/10.1186/s40104-019-0350-9*
18. *Thompson, S. V., Hannon, B. A., An, R., & Holscher, H. D. (2017). Effects of isolated soluble fiber supplementation on body weight, glycemia, and insulinemia in adults with overweight and obesity: a systematic review and meta-analysis of randomized controlled trials. The American journal of clinical nutrition, 106(6), 1514–1528. https://doi.org/10.3945/ajcn.117.163246*
19. *Yao, B., Fang, H., Xu, W., Yan, Y., Xu, H., Liu, Y., Mo, M., Zhang, H., & Zhao, Y. (2014). Dietary fiber intake and risk of type 2 diabetes: a dose-response analysis of prospective studies. European journal of epidemiology, 29(2), 79–88. https://doi.org/10.1007/s10654-013-9876-x*
20. *Yang, J., Wang, H. P., Zhou, L., & Xu, C. F. (2012). Effect of dietary fiber on constipation: a meta analysis. World journal of gastroenterology, 18(48), 7378–7383. https://doi.org/10.3748/wjg.v18.i48.7378*
21. *Anderson, J. W., Allgood, L. D., Lawrence, A., Altringer, L. A., Jerdack, G. R., Hengehold, D. A., & Morel, J. G. (2000). Cholesterol-lowering effects of psyllium intake adjunctive to diet therapy in men and women with hypercholesterolemia: meta-analysis of 8 controlled trials. The American journal of clinical nutrition, 71(2), 472–479. https://doi.org/10.1093/ajcn/71.2.472*
22. *Slavin, J. L., & Lloyd, B. (2012). Health benefits of fruits and vegetables. Advances in nutrition (Bethesda, Md.), 3(4), 506–516. https://doi.org/10.3945/an.112.002154*
23. *Xue, Y., Cui, L., Qi, J., Ojo, O., Du, X., Liu, Y., & Wang, X. (2021). The effect of dietary fiber (oat bran) supplement on blood pressure in patients with essential hypertension: A randomized controlled trial. Nutrition, metabolism, and cardiovascular diseases : NMCD, 31(8), 2458–2470. https://doi.org/10.1016/j.numecd.2021.04.013.*
24. *Jiménez-Cortegana, C., Iglesias, P., Ribalta, J., Vilariño-García, T., Montañez, L., Arrieta, F., Aguilar, M., Durán, S., Obaya, J. C., Becerra, A., Pedro-Botet, J., Sánchez-Margalet, V., & On Behalf Of The Cardiovascular Disease Working Group Of The Spanish Society Of Diabetes Sed (2021). Nutrients and Dietary Approaches in Patients with Type 2 Diabetes Mellitus and Cardiovascular Disease: A Narrative Review. Nutrients, 13(11), 4150. https://doi.org/10.3390/nu13114150*
25. *Baye, K., Guyot, J. P., & Mouquet-Rivier, C. (2017). The unresolved role of dietary fibers on mineral absorption. Critical reviews in food science and nutrition, 57(5), 949–957. https://doi.org/10.1080/10408398.2014.953030*
26. *Wanders, A. J., van den Borne, J. J., de Graaf, C., Hulshof, T., Jonathan, M. C., Kristensen, M., Mars, M., Schols, H. A., & Feskens, E. J. (2011). Effects of dietary fibre on subjective appetite, energy intake and body weight: a systematic review of randomized controlled trials. Obesity reviews : an official journal of the International Association for the Study of Obesity, 12(9), 724–739. https://doi.org/10.1111/j.1467-789X.2011.00895.x*
27. *Noguera-Fernández, N., Candela-González, J., & Orenes-Piñero, E. (2024). Probiotics, Prebiotics, Fecal Microbiota Transplantation, and Dietary Patterns in Inflammatory Bowel Disease. Molecular nutrition & food research, 68(18), e2400429. https://doi.org/10.1002/mnfr.202400429*
28. *Zhang, Y., Belkassem, N., Umanets, A., de Santana-Filho, A. P., Venema, K., Schols, H. A., Penders, J., Jonkers, D., & Abboud, K. Y. (2025). A head-to-head comparison of soluble dietary fibers on microbiota composition and short-chain fatty acid production using the in vitro TIM-2 system. Carbohydrate polymers, 368(Pt 1), 124090. https://doi.org/10.1016/j.carbpol.2025.124090*
29. *Chen, X., Xu, H., Xu, Y., Wu, J., Zou, B., Li, L., Yu, Y., & Hu, T. (2025). The interaction between soluble dietary fiber and bound phenolic from litchi pomace, and its effect on gut microbiota and inflammation. Food chemistry, 491, 145322. https://doi.org/10.1016/j.foodchem.2025.145322*

30. *Gopalakrishnan, V., Saravanan, V., Mahendran, M. I. M. S., Boopathy, V., Vaithianathan, R., Srinivasan, S., & Krishnamurthy, S. (2025). Examining the Role of Virulence Proteins From Pathogenic Anaerobes in the Initiation of Colorectal Cancer. APMIS : acta pathologica, microbiologica, et immunologica Scandinavica, 133(7), e70041. https://doi.org/10.1111/apm.70041*
31. *Barber, T. M., Kabisch, S., Pfeiffer, A. F. H., & Weickert, M. O. (2020). The Health Benefits of Dietary Fibre. Nutrients, 12(10), 3209. https://doi.org/10.3390/nu12103209 32. FDA 2024, https://www.fda.gov/food/nutrition-food-labeling-and-critical-foods/questions-and-answers-dietary-fibre (accessed in 6 June 2025).*
32. *Howlett, J. F., Betteridge, V. A., Champ, M., Craig, S. A., Meheust, A., & Jones, J. M. (2010). The definition of dietary fiber - discussions at the Ninth Vahouny Fiber Symposium: building scientific agreement. Food & nutrition research, 54, 10.3402/fnr.v54i0.5750. https://doi.org/10.3402/fnr.v54i0.5750*
33. *Izumi, S., Isida, K. and Iwamoto, M. (1933). The Mechanism of the Formation of Phytobezoars With Special Reference to the Persimmon Ball. Jap. J. M. Sc., 2: 21*
34. *DeBakey, M. and Oschner,A. (1939). "Bezoars and concretions." Surgery., 5,: 132-160.*
35. *Albostani, A., Kfelati, F., Alsaadi, W., Faraman, R. A., & Farman, A. (2024). Small bowel obstruction due to a meat bolus bezoar: the second case report in literature. Annals of medicine and surgery (2012), 86(2), 1139–1143. https://doi.org/10.1097/MS9.0000000000001633*
36. *Ali, A. Y., Biyikli, A., Mohamed, A. A., Dhaley, A., & İbrahim, İ. G. (2022). Infrequent cause of small bowel obstruction in a child: A lemon. Annals of medicine and surgery (2012), 76, 103488. https://doi.org/10.1016/j.amsu.2022.103488*
37. *Manatakis, D. K., Acheimastos, V., Antonopoulou, M. I., Balalis, D., & Korkolis, D. P. (2019). Gastrointestinal Seed Bezoars: A Systematic Review of Case Reports and Case Series. Cureus, 11(5), e4686. https://doi.org/10.7759/cureus.4686.*
38. *Salada, K. O., & Jacobson, E. (2023). Seeds of Discomfort: An Unusual Case of Pediatric Abdominal Pain. Cureus, 15(12), e50625. https://doi.org/10.7759/cureus.50625*
39. *Gök, A. F. K., Sönmez, R. E., Kantarcı, T. R., Bayraktar, A., Emiroğlu, S., İlhan, M., & Güloğlu, R. (2019). Discussing treatment strategies for acute mechanical intestinal obstruction caused by phytobezoar: A single-center retrospective study. Fitobezoara bağlı akut mekanik intestinal obstrüksiyon için tedavi stratejilerinin tartışılması: Tek merkezli geriye dönük bir çalışma. Ulusal travma ve acil cerrahi dergisi = Turkish journal of trauma & emergency surgery : TJTES, 25(5), 503–509. https://doi.org/10.14744/tjtes.2019.24557*
40. *Luo G. (2025). Clinical applications of carbonated beverages: A case report of large gastric phytobezoar complicated by acute pancreatitis. Medicine, 104(50), e46536. https://doi.org/10.1097/MD.0000000000046536*
41. *Nawab, S., Shah, M. A., Nambiar, R., Longi, A. A., & Alexander, G. (2026). Phytobezoar-Induced Jejunal Obstruction in an Elderly Male With Multiple Comorbidities. Cureus, 18(3), e105285. https://doi.org/10.7759/cureus.105285*
42. *Khan, H., Shah, A., Zindani, S., & Rahman, K. (2025). S4718 Phytobezoar-Related GI Hemorrhage in the Elderly: An Underrecognized Cause of Upper GI Bleeding. Official journal of the American College of Gastroenterology| ACG, 120(10S2), S1008.*
43. *Doughty, M., Alookaran, J., Rhoads, M., & Dahm, P. H. (2021). Restrictive Lung Pattern: A Rare Complication of a Phytobezoar. JPGN reports, 2(2), e056. https://doi.org/10.1097/PG9.0000000000000056*
44. *Parr, A. R., Horn, C. B., & Freeman, C. A. (2020). Jejunal phytobezoar complicated by small bowel perforation. The journal of trauma and acute care surgery, 88(2), e101–e103. https://doi.org/10.1097/TA.0000000000002531*
45. *Khan, H., Mashood, M., Ahmad, W. and Iqbal, M.O., Proximal Ileal Perforation Secondary due to Phytobezoar in Elderly Patient. World Journal of Research and Review (WJRR), 2024, 19(6), 1-3. https://doi.org/10.31871/WJRR.19.6.28*
46. *Özsoy, Z., & Okan, İ. (2018). Treatment of phytobezoars: Tailoring management to diverse presentations. Turkish journal of surgery, 34(1), 33–37. https://doi.org/10.5152/turkjsurg.2017.3770*
47. *https://pubmed.ncbi.nlm.nih.gov/*
48. *https://www.nlm.nih.gov/medline/medline_home.html*
49. *50.https://www.webofscience.com/wos/woscc/smart-search*
50. *https://scholar.google.com/*
51. *Delgado Galan, M., & Rabago, L. R. (2024). Has Coca-Cola treatment become the first-line*

*therapy for gastric bezoars, both in general and specifically for western countries?. World journal of gastrointestinal endoscopy, 16(5), 237–243. https://doi.org/10.4253/wjge.v16.i5.237*

52. *Mejri, A., & Trigui, E. (2021). Phytobezoar: A train can hide another. International journal of surgery case reports, 81, 105814. https://doi.org/10.1016/j.ijscr.2021.105814*
53. *Fazio, R., Shah, P., Soe, E., Iswara, K., & Chen, I. (2016). Gastric Bezoar Causing Massive Gastric Distention and Functional Outlet Obstruction in a Patient With Bulimia Nervosa. Journal Of Medical Cases, 7(7), 312-314. doi: https://doi.org/10.14740/jmc2560w*
54. *Andrus, C. H., & Ponsky, J. L. (1988). Bezoars: classification, pathophysiology, and treatment. The American journal of gastroenterology, 83(5), 476–478.*
55. *Mahir, S., Salih, A. M., Ahmed, O. F., Kakamad, F. H., Salih, R. Q., Mohammed, S. H., Usf, D. C., Hassan, H. A., & Abdulla, F. (2020). Giant phytobezoar; an unusual cause of gastric outlet obstruction: A case report with literature review. International journal of surgery case reports, 67, 154–156. https://doi.org/10.1016/j.ijscr.2020.02.012*
56. *Buchholz, R. R., & Haisten, A. S. (1972). Phytobezoars following gastric surgery for duodenal ulcer. The Surgical clinics of North America, 52(2), 341–352. https://doi.org/10.1016/s0039-6109(16)39686-4*
57. *Erzurumlu, K., Malazgirt, Z., Bektas, A., Dervisoglu, A., Polat, C., Senyurek, G., Yetim, I., & Ozkan, K. (2005). Gastrointestinal bezoars: a retrospective analysis of 34 cases. World journal of gastroenterology, 11(12), 1813–1817. https://doi.org/10.3748/wjg.v11.i12.1813*
58. *Vellar, D. J., Vellar, I. D., Pucius, R., & Steedman, P. K. (1986). Phytobezoars--an overlooked cause of small bowel obstruction following vagotomy and drainage operations for duodenal ulcer. The Australian and New Zealand journal of surgery, 56(8), 635–638. https://doi.org/10.1111/j.1445-2197.1986.tb04519.x*
59. *Krausz, M. M., Moriel, E. Z., Ayalon, A., Pode, D., & Durst, A. L. (1986). Surgical aspects of gastrointestinal persimmon phytobezoar treatment. American journal of surgery, 152(5), 526–530. https://doi.org/10.1016/0002-9610(86)90221-7*
60. *Zheng, Y. X., Prasoon, P., Chen, Y., Hu, L., & Chen, L. (2014). ''Sandwich" treatment for diospyrobezoar intestinal obstruction: a case report. World journal of gastroenterology, 20(48), 18503–18506. https://doi.org/10.3748/wjg.v20.i48.18503*
61. *Shiotsuki, K., Imai, K., & Hotta, K. (2019). Small bowel obstruction by massive impacted diospyrobezoars relieved by endoscopic lithotripsy with a polypectomy snare using double balloon endoscopy. Digestive endoscopy : official journal of the Japan Gastroenterological Endoscopy Society, 31(6), e111–e112. https://doi.org/10.1111/den.13490.*
62. *63. USDA: https://fdc.nal.usda.gov/food-search?type=SR%20Legacy&query=persimmon (accessed in 20/6/2025)*
63. *Marchese, S., Bertucci, B., Manti, F., Berritto, D., Roperto, A. G., & Tamburrini, S. (2015). RECTAL IMPACTION DUE TO PRICKLY PEAR SEEDS BEZOAR: A CASE REPORT. Journal of biological regulators and homeostatic agents, 29(3), 707–711.*
64. *Su, W., Chen, J., Zhao, Y., Xu, P., Yuan, J., Luo, C., Liu, J., & Wang, B. (2025). Diagnosis and treatment of giant colonic fecalith in a child: a case report. Frontiers in pediatrics, 13, 1598807. https://doi.org/10.3389/fped.2025.1598807.*
65. *Gentile, M., Vergara, L., Schiavone, V., Cestaro, G., & Sivero, L. (2022). Harry Potter's Occlusion: Report of a Case of Pumpkin Seed Bezoar Rectal Impact. Frontiers in surgery, 9, 902701. https://doi.org/10.3389/fsurg.2022.902701.*
66. *Fernando, A. R., Bulathsinghela, R., & Samarasekera, D. N. (2017). An unusual cause of small bowel obstruction due to an ingested mango seed: a case report. BMC research notes, 10(1), 549. https://doi.org/10.1186/s13104-017-2875-3.*
67. *Ozcan, U. A., Yilmaz, S., Akansel, S., Cizmeli, M. O., & Ertem, M. (2007). An unusual cause of small bowel obstruction: CT diagnosis of dried apricots. Emergency radiology, 14(6), 417–419. https://doi.org/10.1007/s10140-007-0625-x.*
68. *Atila, K., Güler, S., Bora, S., & Gülay, H. (2011). An unusual cause of small bowel perforation: apricot pit. Ulusal travma ve acil cerrahi dergisi = Turkish journal of trauma & emergency surgery : TJTES, 17(3), 286–288.*
69. *Al-Najjar, M., & Arthur, T. (2013). The nectarine pit as a cause for small bowel obstruction and perforation: a case report. Case reports in surgery, 2013, 902943. https://doi.org/10.1155/2013/902943.*

70. *Geerts, B., Verstraeten, T., Vanslambrouck, K., Vandenhaute, S., Dillemans, B., & Coenegrachts, K. (2015). Unusual Cause of Small Bowel Obstruction Caused by a Prune Pit. Journal of the Belgian Society of Radiology, 99(1), 95–96. https://doi.org/10.5334/jbr-btr.838.*
71. *Okagawa, Y., Takada, K., Arihara, Y., & Kato, J. (2017). Nihon Shokakibyo Gakkai zasshi = The Japanese journal of gastro-enterology, 114(10), 1830–1835. https://doi.org/10.11405/nisshoshi.114.1830*
72. *Ravindra, R. K., Das, A., Chew, G. L., & Daniel, E. (2019). Small bowel obstruction with multiple perforations post chestnut ingestion. BMJ case reports, 12(2), bcr-2018-227999. https://doi.org/10.1136/bcr-2018-227999.*
73. *FDA (Food and Drug Administration), HHS. (2007). Laxative drug products for over-the-counter human use; psyllium ingredients in granular dosage forms. Final rule. Fed Regist, 72, no. 60 (2007): 14669-74. PMID: 17450664. https://www.govinfo.gov/content/pkg/FR-2007-03-29/pdf/E7-5740.pdf*
74. *Hefny, A. F., Ayad, A. Z., Matev, N., & Bashir, M. O. (2018). Intestinal obstruction caused by a laxative drug (Psyllium): A case report and review of the literature. International journal of surgery case reports, 52, 59–62. https://doi.org/10.1016/j.ijscr.2018.10.001.*
75. *Salguero Molpeceres, O., Seijas Ruiz-Coello, M. C., Hernández Núñez, J., Caballos Villar, D. (2003). "Obstrucción esofágica por la fibra dietética Plantago ovata, una complicación prevenible mediante la información" [Esophageal obstruction caused by dietary fibre from Plantago ovata, a complication preventable by adequate information]. Gastroenterol Hepatol., 26, no. 4: 248-50. PMID: 12681118.*
76. *Agha, F. P., Nostrant, T. T., & Fiddian-Green, R. G. (1984). "Giant colonic bezoar:" a medication bezoar due to psyllium seed husks. The American journal of gastroenterology, 79(4), 319–321.*
77. *Shin, S., Kim, J. H., Mun, Y. H., & Chung, H. S. (2022). Acute esophageal obstruction after ingestion of psyllium seed husk powder: A case report. World journal of clinical cases, 10(7), 2336–2340. https://doi.org/10.12998/wjcc.v10.i7.2336.*
78. *Rawl,R. and Browne, L. (2014). Watch It Grow: Esophageal Impaction With Chia Seeds: 833. Am J of Gastroenterol, 109, 1: S244-S245,*
79. *Yeoh S. W. (2017). Esophageal bezoar due to karaya gum granules used as a laxative. Clinical journal of gastroenterology, 10(5), 437–441. https://doi.org/10.1007/s12328-017-0764-x*
80. *Schlag, S., Liese, J., & Veldhoen, S. (2021). Akutes Abdomen mit galligem Erbrechen – Ein Leinsamen-Bezoar als ungewöhnliche Ursache eines obstruktiven Ileus bei einer Adoleszenten mit Anorexie [Acute abdomen with Bilious Vomiting - A Linseed Bezoar as a Rare Cause of an Obstructive Ileus in an Anorexic Adolescent]. Klinische Padiatrie, 233(2), 72–74. https://doi.org/10.1055/a-1170-7435.*
81. *Kang, J. Y., & Doe, W. F. (1979). Unprocessed bran causing intestinal obstruction. British medical journal, 1(6173), 1249–1250. https://doi.org/10.1136/bmj.1.6173.1249-a*
82. *Allen-Mersh, T., & De Jode, L. R. (1982). Is bran useful in diverticular disease?. British medical journal (Clinical research ed.), 284(6317), 740. https://doi.org/10.1136/bmj.284.6317.740*
83. *McClurken, J. B., & Carp, N. Z. (1988). Bran-induced small-intestinal obstruction in a patient with no history of abdominal operation. Archives of surgery (Chicago, Ill. : 1960), 123(1), 98–100. https://doi.org/10.1001/archsurg.1988.01400250108021.*
84. *Rosario, P. G., Gerst, P. H., Prakash, K., & Albu, E. (1990). Dentureless distention: oat bran bezoars cause obstruction. Journal of the American Geriatrics Society, 38(5), 608. https://doi.org/10.1111/j.1532-5415.1990.tb02419.x*
85. *Piotto, L., & Gent, R. (2005). Dried apricots: an unusual cause of bowel obstruction. Pediatric radiology, 35(12), 1224–1226. https://doi.org/10.1007/s00247-005-1552-1*
86. *Ooi, S., & Hong, K. (2015). Small bowel obstruction caused by dried apple. International journal of surgery case reports, 10, 154–157. https://doi.org/10.1016/j.ijscr.2015.03.038.*
87. *Oka, A., Amano, Y., Uchida, Y., Kagawa, K., Takatori, K., Kitajima, N., Sonoyama, H., Tada, Y., Kusunoki, R., Fukuba, N., Oshima, N., Moriyama, I., Yuki, T., Kawashima, K., Ishihara, S., & Kinoshita, Y. (2013). Small bowel obstruction and gastric ulceration resulting from rice cake ingestion-computed tomography diagnosis in eight patients. Nihon Shokakibyo Gakkai Zasshi= The Japanese Journal of Gastroenterology, 110(10), 1804-1813.*
88. *Oka, A., Ishihara, S., Mikami, H., Sonoyama, H., Mishiro, T., Tobita, H., Kawashima, K., Miyake, T., Ishimura, N., Furuta, K., Kinoshita, Y., & Nishina, M. (2019). Retained Rice Cake: A Unique Upper Gastrointestinal*

*Foreign Body: Case Report and a Literature Review. Internal medicine (Tokyo, Japan), 58(17), 2485–2494. https://doi.org/10.2169/internalmedicine.2760-19*

89. *Chao, H. C., Chang, K. W., & Wang, C. J. (2012). Intestinal obstruction caused by potato bezoar in infancy: a report of three cases. Pediatrics and neonatology, 53(2), 151–153. https://doi.org/10.1016/j.pedneo.2012.01.015*
90. *Sugimoto, S., Shimono, T., Takeshita, T., Yamamoto, A., Shindo, D., & Miki, Y. (2019). Clinical and CT findings of small bowel obstruction caused by rice cakes in comparison with bezoars. Japanese journal of radiology, 37(4), 301–307. https://doi.org/10.1007/s11604-019-00811-9*
91. *Katsinelos, P., Pilpilidis, I., Chatzimavroudis, G., Katsinelos, T., Lazaraki, G., Fasoulas, K., Zavos, C., & Kountouras, J. (2009). Huge gastric bezoar caused by honeycomb, an unusual complication of health faddism: a case report. Cases journal, 2, 7077. https://doi.org/10.1186/1757-1626-2-7077*
92. *Hsu, H. H., Grove, W. E., Mindulzun, R., & Knauer, C. M. (1992). Gastric bezoar caused by lecithin: an unusual complication of health faddism. The American journal of gastroenterology, 87(6), 794–796.*
93. *Beyene, B., Mohammadsani, B., Abdlhadi, M., Getachew, K., & Ahmedin, H. (2022). Khat induced bezoar; a rare cause of small bowel obstruction:a case report. International journal of surgery case reports, 91, 106811. https://doi.org/10.1016/j.ijscr.2022.106811*
94. *Huang, Y., Meng, J., & Zhang, S. (2025). Small intestinal obstruction caused by the gastric bezoars in elderly diabetic patients: A case report. International journal of surgery case reports, 127, 110761. https://doi.org/10.1016/j.ijscr.2024.110761*
95. *Lim, C. H., Lim, C. J., Yao, C. T., & Chang, C. C. (2025). Novel approach to managing two enormous bezoars with successive snare-tip electrocautery: A case report. World journal of gastrointestinal endoscopy, 17(1), 102185. https://doi.org/10.4253/wjge.v17.i1.102185*
96. *Patiño-Araujo, B., Salgado-Báez, M. E., Villavicencio-Logroño, G. P., Ayora-Pérez, M. J., Zapata-Jaramillo, G. S., & Salgado-Macías, N. (2025). Early small bowel obstruction by phytobezoar related to zopiclone induced amnesia after revisional Roux-en-Y gastric bypass: a case report. Journal of surgical case reports, 2025(4), rjaf202. https://doi.org/10.1093/jscr/rjaf202*
97. *Boopathy, V., & Palaniswamy, M. (2014). A giant complex rectosigmoid bezoar with multiple foreign bodies: a first of its kind. BMJ case reports, 2014, bcr2014207573. https://doi.org/10.1136/bcr-2014-207573.*
98. *Paschos, K. A., & Chatzigeorgiadis, A. (2019). Pathophysiological and clinical aspects of the diagnosis and treatment of bezoars. Annals of gastroenterology, 32(3), 224–232. https://doi.org/10.20524/aog.2019.0370*
99. *Liu, B., & Kuang, S. (2024). Gastric Bezoars: A Retrospective Analysis of 44 Cases. Medical science monitor : international medical journal of experimental and clinical research, 30, e945377. https://doi.org/10.12659/MSM.945377.*
100. *Hemmasi, G., Zanganeh, E., Hosseini, S. A., Alimadadi, M., Ahmadi, A., Hajiebrahimi, S., & Seyyedmajidi, M. (2021). Risk factors, endoscopic findings, and treatments in upper gastrointestinal bezoars: multi-center experience in Iran. Gastroenterology and hepatology from bed to bench, 14(2), 160–164.*
101. *De Cesare, A., Fiori, E., Bononi, M., & Ferraro, D. (2015). Phytobezoar-induced small bowel obstruction associated with a concomitant gastric phytobezoar and ulcer in an elderly woman. Annali italiani di chirurgia, 86(1), 70–77*
102. *Khan, S., Jiang, K., Zhu, L. P., Khan, I. A., Ullah, K., Khan, S., Chen, X., & Wang, B. M. (2019). Upper Gastrointestinal Manifestation of Bezoars and the Etiological Factors: A Literature Review. Gastroenterology research and practice, 2019, 5698532. https://doi.org/10.1155/2019/5698532*
103. *Ugenti, I., Travaglio, E., Lagouvardou, E., Caputi Iambrenghi, O., & Martines, G. (2017). Successful endoscopic treatment of gastric phytobezoar: A case report. International journal of surgery case reports, 37, 45–47. https://doi.org/10.1016/j.ijscr.2017.06.015*
104. *Aydin, I., Sengul, I., & Sengul, D. (2022). Phytobezoar: An Unusual Condition Leading to Small Bowel Obstruction. Cureus, 14(4), e23885. https://doi.org/10.7759/cureus.23885*
105. *Chineka, R., Mukosera, T., Dzvanga, N., & Gwanda, B. (2021). Case report-An unusual diagnosis of a gastric phytobezoar in a middle-aged woman. Clinical case reports, 9(5), e04077. https://doi.org/10.1002/ccr3.4077*
106. *107.Asefa, H. Y., Wassie, M. T., Assefa, B. C., Sefene, B. S., Yilma, Y. F., & Dereje, W. M. (2025). Phytobezoar causing small bowel obstruction in a 4-year-old girl: a case report. Journal of Pediatric Surgery Case Reports, 103138.*

**107. S1. Appendix:** list of reviewed phytobezoar cases.

**Table 1: List for phytobezoar cases associated with previous surgery.**

| Gender | Age (years) | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|
| Female | 51 | laparoscopic cholecystectomy | Diabetes mellitus | [1] |
| Male | 65 | duodenal ulcer surgery | anamnesis of recurrent ileus | [2] |
| Male | NA | laparoscopic sleeve gastrectomy (SG) | | [3] |
| Male | NA | laparoscopic sleeve gastrectomy (SG) | | [4] |
| Male | 55 | Stapled gastroplasty | Nephrolithiasis | [5] |
| Male | NA | laparoscopic sleeve gastrectomy (SG) | | [6] |
| Male | NA | laparoscopic sleeve gastrectomy (SG) | | [7] |
| Female | 33 | She had a laparo- scopic Roux-en-Y gastric bypass (LRYGB) for morbid obesity | Obese, Body Mass index: 36. | [8] |
| 150 cases, gender is not specified | NA | All have had previous gastrectomy for duodenal ulcer. | High fiber foods | [9] |
| Female | 63 | RYGB | | [10] |
| 13 cases (7 males and 6 females) | Range: 44-79 | 11 patients had previous ulcer surgery, 8 truncal vagotomy with pyloroplasty and 3 partial gastrectomy | Ingestion of persimmon, grapefruits and dried plumes | [11] |
| Female | 68 | subtotal gastrectomy due to gastric cancer | dyslipidemia, supraventricular paroxysmal tachycardia anddepression | [12] |
| Male | 75 | surgical resction of the sigmoid colon for diverticulitis | he ingested approximately 60 g Oat bran daily | [13] |
| Male | 48 | Laparoscopic adjustable gastric banding (LAGB) | | [14] |
| Female, | 50 | Laparoscopic sleeve astrectomy (in 2014) Roux-en-Y gastric bypass (RYGB) (in 2017) | Poor mastication | [15] |
| Male | 66 | Billroth II hemigastrectomy associated with truncal vagotomy to treat a chronic duodenal ulcer | Diabetes | [16] |
| Female (pregnant) | 33 | RYGB | | [8] |
| Female | 59 | laparoscopic adjustable gastric banding (LAGB) for morbid obesity | Obesity and related issues. | [17] |
| Female | 75 | a subtotal gastrectomy and Billroth II gastrojejunostomy were performed. | NA | [18] |
| Female | 63 | laparoscopic Roux-en-Y gastric bypass surgery | morbid obese | [19] |
| 18 Males/14 Females (1.42/1). | 32 cases, (range 17-78) years; Mean age was 53.73±15.56 | (19 of them have previous surgery.) 13 have had another predisposing factors. | Peptic ulcera (n = 4, 11.76%), diabetes mellitus (n = 4, 11.76%), cherrylaurel (Prunus Buricerasus) or persimmon (Diospyrus Lotus) intakes (n = 6, 17.64%), mental retardation-trichotillomania (n = 2, 5.88%), high dose intake of H2 receptor antagonists (n = 1, 2.94%) were also found in patients' history. | [20] |

| Gender | Age (years) | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|
| Male | 47 | Yes | Raw stinging nettle | [21] |
| Female | 86 | appendectomy | | [22] |
| Female | 64 | Laparoscopic Roux-en-Y gastric bypass | | [23] |
| Female | 51 | RYGB | | [24] |
| Male | 73 | Billroth I partial gastrectomy for gastric cancer. | | [25] |
| Female | 58 | Laparoscopic adjustable gastric banding | | [25] |
| 43 cases (23 males and 20 females ) | Range: 18-93 | 8 cases, (18.6%) Gastrectomy | Diabetes, dental problems, Stomach diseases accounted for 25.5%, 43.6%, 69.7% respectively. | [26] |
| 95 cases of 103 (52 Males and 43 Females) | Mean Age : 60.5 ± 11.3) years | gastric surgery (25.2%), | risk factors were history of gastric surgery (25.2%), diabetes mellitus (21.3%), hypothyroidism (15.5%), trichophagia (5.8%), and anxiety disorders (2.9%), respectively | [27] |
| Female | 55 | RYGB | | [28] |
| Female | 69 | open appendicectomy | Mango seed | [29] |
| Female | 66 | Appendectomy | Liver cirrhosis | [30] |
| Male | 59 | Subtotal gastrectomy | | [30] |
| Male | 70 | Ivor-Lewis operation | Lung cancer | [30] |
| Male | 55 | Primary repair for duodenal ulcer | | [30] |
| Male | 81 | laparoscopic cholecystectomy, surgical repair of bilateral inguinal hernias with mesh placement | acute myocardial infarction, hypertension, surgery related ileus. | [31] |
| (78 male and 35 female) | NA | 106 patients have had previous surgery for duodenal ulcer. | 103 of 113 cases, reported persimmon ingestion prior to GI obstruction. | [32] |
| Female | 19 | No | Anorexia nervosa , Binge-purge , Hematemesis | [33] |
| (20 males, 24 females) | (55.36±15.17) | 7 cases out of 44 patients, have had previous surgeries | Relatively older patients (55.36±15.17), Surgery (7 cases), Chronic diseases (13 cases) | [34] |
| 90 Females & 63 M ales including (29 girls + 54 Boys) | Age range (2-80 Y | Only 10 surgery cases | 18 cases (1 child, 17 adults) have predisposing factors: gastrointestinal strictures (10/153, 6.5%), diabetes (4/153, 2.6%) and neuropsychiatric disease (4/153, 2.6%). | [35] |
| Male | | laparoscopic sleeve gastrectomy (SG) | | [36] |
| 5 Male, 9 Female | 52 -82 Y | 4 appendictomy (2F+2M), 1 hysterctomy, (1, M) gasterectomy, 1 ileus. +3 cases female cesarean resections | All patients had ingested rice cake by swallowing without chewing. | [37] |
| Female | 93 | distal gastrectomy and laparoscopic cholecystectomy | Ingestion of Kaky fruit | [38] |
| 5 Male, 3 Female | 62, 63, 68, 76, 85 (M); 56, 65, 83 (F) | abdominal syrgeries, 2 cases (1 m and 1F) | 5 have dentures (3M+2F), 1 quick feeder, 1had operation, 1 operation and quick feeder (F) | [39] |
| Female | 34Y | LAGB | | [40] |
| Female | 54Y | RYGB | | [41] |

| Gender | Age (years) | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|
| 4 Females, 5 Males | 52-83 (mean age : 67.3), | 5/9 had gastric surgery, | Aging, gastric surgery, Poor mastication, medications DM., hypertention + hyperlipidemia, Abdomino-perineal resection 1/9Unremarkable case (67 Y. old) | [42] |
| Female | 51Y | RYGB | | [43] |
| Hispanic Male | 13Y., | Appendectomy, 6 year ago. | large amount of white fibrous material of unclear etiology | [44] |
| Female | 57 Y | RYGB | | [45] |
| Female | 34 Y, | laparoscopic Roux-en-Y gastric bypass (RYGB) due to class III morbid obesity | Morbid obesity | [46] |
| Male | 57 Y. | Gastric laparotomy | Hypertension | [47] |
| Male | 78 | Gasterctomy | Diospyrobezoar | [48] |
| Male | 34 Y, | RYGB | | [49] |
| Female | 49 Y | RYGB | | [50] |
| Male | 57 Y, | RYGB | | [51] |
| Female | 46 Y | RYGB | | [51] |
| Male, | 57 Y, | Laparoscopic Roux-en-Y gastric bypass | Morbid obesity, BMI of 44, hypertension, obstructive sleep apnea, and ventral hernia due to previous open cholecystectomy underwent laparoscopic gastric bypass | [51] |
| Female | 46 Y | Laparoscopic Roux-en-Y gastric bypass | Morbid obesity, BMI of 44, hypertension, obstructive sleep apnea, and ventral hernia due to previous open cholecystectomy underwent laparoscopic gastric bypass | [51] |
| Male | 70s Y. | open cholecystectomy | Rapid consumption an expiring bag of peanuts, without perfect chewing. | [52] |
| Female | 66 | pylorus-preserving gastrectomy | large amount of persimmon fruits | [53] |
| 9Male/ 8Female | (44-80) | Appendicectomy (3 cases), Hysterectomy (4 cases) | Gulping habit (2case), toothless (1case), | [54] |
| Female | F, 86 Y., solid food bezoar | Cholecystectomy surgery | Ingestion of solid food bezoar | [55] |
| Japanese male, | 56 Y. | pigtail-shaped PEG-J (percutaneous endoscobic transgastric jejunectomy) tube | Levodopa-carbidopa intestinal gel (LCIG) is an established device-aided therapy for patients with advanced Parkinson's disease (PD) | [56] |
| (7 Males, & 4 Females), | Age ranged between 42-74 Y olds , | pigtail-shaped PEG-J (percutaneous endoscobic transgastric jejunectomy) tube | levodopa-carbidopa intestinal gel (LCIG) is an established device-aided therapy for patients with advanced Parkinson's disease (PD) | [56] |
| 6 Males, 3 Females | 50-65 (males), 36, 56, 69 (females) | All have had previous vagotomy and drainage procedures for duodenal ulcer. | Not mentioned | [57] |
| Female | 47Y | RYGB | | [58] |
| Female | 37Y | LAGB | | [58] |
| Female | 42Y | LAGB | | [59] |
| 51 cases 27 Males : 24 Females | Median Age 62.9 (range: 27-89) years | Previous gastric surgery (n = 16, 31.4%) was the most common predisposing risk factor | Other predisposing factors: diabetes, hypertension, peptic ulcer, persimmon fruit, brown see weeds, and others | [60] |

| Gender | Age (years) | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|
| Female | 21 Y | RYGB | | [61] |
| Male | A 47-years | Yes | underwent a cholecystectomy 10 years ago<br>the patient admitted to consuming a diet high in fibrous vegetables<br>proton pump inhibitors | [62] |
| Female | 59-year | Yes | Roux-en-Y gastric bypass (RYGB) | [63] |
| Male | 67-Year | Yes | Prostatic cancer - radiotherapy | [147] |
| Female | 45-Year | Yes, | | [144] |
| Female | 40-Year | Yes, | 2 times Cholecystectomy | [137] |

**Table 2: List for diospyrobezoar (persimmon) cases.**

| Gender | Age | previous surgery | Risk factors & medical history | Ref. |
|---|---|---|---|---|
| 96 cases (22 females , 74Males) | NA | no | NA | [64] |
| 20 male and 1Female | 40-76 Y | no | persimmon fruits | [65] |
| Male | 73 | no | persimmon fruits | [66] |
| (78 male and 35 female) | 113 cases | 106 patients have had previous surgery for duodenal ulcer. | 103 of 113 cases, reported persimmon ingestion prior to GI obstruction. | [32] |
| Male, | 35 Y., | No | large quantity of unripe persimmons | [67] |
| 49 Males +10 Females | 2 -78 Y. (median 56) | Not mentioned | Eating persimmon's fruits | [68] |
| 51 cases 27 Males : 24 Females | Median Age 62.9 (range: 27-89) years | Previous gastric surgery (n = 16, 31.4%) was the most common predisposing risk factor | Other predisposing factors: diabetes, hypertension,peptic ulcer, persimmon fruit (11 cases), brown see weeds ( 4cases), and others | [60] |
| Male | 27 | no | large amount of persimmon fruits | [69] |
| Girl | 5-Years | No | Persimmon | [142] |
| Female | 88-Years | No | Persimmon | [139] |
| Female | 54-Years | No | Persimmon | [145] |

**Table 3- list for seed bezoar cases**

| Gender | Age (years) | phytobezoar Type | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Male | 78 | Giant granadilla's seed bezoar (Passiflora quadrangularis) | no | essential hypertension and prostate benign hyperplasia | [70] |
| 2 Males | 6 &10 | unshelled sunflower seeds and areca nuts. | No. | unshelled sunflower seeds and areca nuts. | [71] |
| Male | 1.7 | Male, 1.7 Y., seed shells | No | Hard fibrous plant material, (seed shells). | [72] |

| Gender | Age (years) | phytobezoar Type | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| 16 Male/14 Female | Range: 19 - 79 | prickly pear seeds (12 cases), watermelon seeds in (10), sunflower seeds (6), popcorn kernels (1) and pomegranate (1). | not specified | (5 cases ) have predisposing factors: neurologic, psychiatric, or anal disorder, as well as previous surgery | [73] |
| 18 Males/14 females (1.29/1). | 32 cases, (range 17-78) years; Mean age was 53.73±15.56 | 32 cases phytobezoar , | (19 of them have previous surgery.) 13 have had another predisposing factors. | Peptic ulcera (n = 4, 11.76%), diabetes mellitus (n = 4, 11.76%), cherrylaurel (Prunus Buricerasus) or persimmon (Diospyrus Lotus) intakes (n = 6, 17.64%), mental retardation-trichotillomania (n = 2, 5.88%), high dose intake of H2 receptor antagonists (n = 1, 2.94%) were also found in patients' history. | [20] |
| Female | 21 | phytobezoar (Food matter) | No | Bulimia nervosa Binge eating episodes | [74] |
| 78 cases | NA | psyllium bezoar | No | | [75] |
| Male | 40 | Mango seed bezoar | No | The patient was under the influence of alcohol, accidental ingestion of mango seed occurred., | [76] |
| Male, Unemployed , | 50 Y | pumpkin seed bezoar. | No | -pumpkin seeds (about 400 gr.) with their shell. | [77] |
| Male | 4y | undisegisted rhubarb | No | undigested rhubarb | [78] |
| 43 cases (23 males and 20 females ) | Range: 18-93 | 43 cases; Suffered from intestinal obstruction due to phytobezoars (40.5% persimmon's bezoars; and 43.6 % gauava's bezoars) | 8 cases, (18.6%) Gastrectomy | Diabetes, dental problems, Stomach diseases accounted for 25.5%, 43.6%, 69.7% respectively. | [26] |
| Male | 64 | Man, 64 Y, phytobezoar due to unshelled sunflower seeds. | No | No | [79] |
| 2 Males | 13 & 6 years | unshelled sunflower seeds | No. | Have had consumed unshelled sunflower seeds | [80] |
| 2 Males | 8 &10 Y, | 2 boys, 8 &10 Y, phytobezoar due to large amount of unshelled sunflower seeds | No | N.A. | [81] |
| Male | 72 | Male, 72 Y., Sunflower-fruits | No | large amount of fried, unshelled sunflower seeds | [82] |
| Male | 12 | Rectal bezoar sunflower seeds | No | 200 grams of salted sunflower seeds with shells | [83] |
| 16 Males/14 Females | 19 - 79 | unshelled watermelon seeds (10 cases), | not determined | (5 cases ) have predisposing factors: neurologic, psychiatric, or anal disorder, as well as previous surgery | [73] |
| Male | 7 | seed bezoar watermelon | No | Unshelled watermelon seeds | [84] |
| Female | 76 | prickly pear seeds | No | diabetes, controlled only by diet, 40 prickly pear fruits with their seeds | [85] |

| Gender | Age (years) | phytobezoar Type | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Male | 48 | a huge vegetable materials | No | hypertension and kidney failure , high fiber diet | [86] |
| Male | 21 | psyllium seed husk bezoar | No | Psyllium husk ingestion without andequate amount of fluid | [87] |
| 95 cases of 103 (52 Males and 43 Females) | Mean Age ( 60.5±11.3) years | 95 cases phytobezoar | gastric surgery (25.2%), | risk factors were history of gastric surgery (25.2%), diabetes mellitus (21.3%), hypothyroidism (15.5%), trichophagia (5.8%), and anxiety disorders (2.9%), respectively | [27] |
| Female, | 43 Y, | 43 Y, Female, (olive seeds phytobezoar) | No | • proton pump inhibitors, due to duodenal ulcer<br>• pyloric stenosis | [88] |
| Male | 49 | Jujubes | No | Habitual jujubes ingestion | [89] |
| 90 Females & 63 M ales including (29 girls + 54 Boys) | Age range (2-80 Y.). 54Boy+29 girl | 153 cases of seed bezoar<br>55 cases (unshelled watermelon seeds),<br>30 unshelled sunflower seeds, 28 prickly pears, 11 willed banana seeds.5 unshelled pumpkin seeds, 4 pomegranate seeds. 2 cases have been reported for each of Date, olive pits, tangerine, lupin and popcorn seeds. While 1 case for granadilla, sesame, Baccaurea macrocarpa, jaboticaba, peanuts, lentil, cherry pits, box myrtle, mango seeds and freekeh grains. | No | 18 cases (1 child, 17 adults) have adults) have predisposing factors: gastrointestinal strictures (10/153, 6.5%), diabetes (4/153, 2.6%) and neuropsychiatric disease (4/153, 2.6%). | [35] |
| Male | 79 | Sesame seed bezoar. | Yes | He underwent a sigmoid colectomy, with histology showing diverticular disease | [90] |
| Female, (housewife) . | 41 Y., | Phytobezoar formed of undigested vegetable matter (peas), fibrous material and decomposed food residue. | No surgery | No (may be poor physical exercise leads to GI laziness) | [91] |
| Boy | 7 years | Ingestion of a large amount of unshelled broad beans. | No | No | [92] |
| 4 Male cases | (15, 16, 22 &18) | Phytobezoar from jungle banana seeds | No | Ingestion of large quantity of jungle banana with seeds. | [93] |
| Girl | 2 years | Excessive consumption of peas | No | No | [94] |
| Female | 54 | Fresh hawthorns 200-250 g every day for a year | No | fresh hawthorns 200-250 g every day for a year | [95] |
| Female | 86 | Phytobezoar due to apricot pits | No | she suffered from difficulties in chewing foods due to her denture, | [96] |
| Female | 73 | Phytobezoar due to apricot pits | No | She swallowed the pit accidently | [97] |
| Male | 67 | Nectarine pit | No | he swallowed the pit accidently | [98] |
| Male | 86 | Prune pit bezoar | Yes | He has had previous appendectomy | [22] |
| Boy | 10-years | Seed besoar | No | | [138] |
| Woman | 60-years | Seed bezoar | No | | [138] |

| Gender | Age (years) | phytobezoar Type | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Man | 25-Years | Guava bezoar | No | | [148] |
| Female | 45-Years | Coconut bezoar | yes | | [144] |

**Table 4: list for phytobezoar cases due to bulk producers**

| Gender | Age (years) | phytobezoar Type | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Male | 50 | bran | No | (two large Bowls of bran cereal with milk) | [99] |
| Female | 84 | Oat bezoar | No | dentureless, poor mustication, High fiber diet | [100] |
| Male | 39 | Chia seed | No | asthma and seasonal allergies | [101] |
| Female | 41 | psyllium seed bezoar | No | she swallowed dry psyllium seed | [102] |
| Female | 14 | Linseed bezoar | No | ingestion of great amount of linseed | [103] |
| Male | 76 | psyllium seed bezoar | No | ingestion of psyllium seed husk powder, Parkinson's disease | [104] |
| Man | 66 | karaya gum | No | This occurred after ingesting several tablespoons of a granular stool bulking laxative containing 62% karaya gum for chronic constipation, prior to reclining to sleep. He did not drink any liquids after ingesting these granules. | [105] |
| 78 cases | | phytobezoars or GIO due to ingestion of laxative drug formed mainly of psyllium seeds or psyllium seed husks. | No | | [75] |
| Male | 21 | psyllium seed husks | No | He did not drink the required amount of fluids because he wasfasting. | [87] |
| Female | 74 | psyllium seed | No | | [106] |
| Female | 59 | Agiolax® (psyllium seed, psyllium husks, senna pods) | No | body mass index (BMI) of 37.9 kg/m and multiple co-morbidities (hypertension, diabetes, hyperuricemia, hyperlipidemia, depression and osteoarthrosis of low back) | [107] |
| Male | 91 | Sterculia gums | No | after mistakenly swallowing a tablespoon of sterculia granules | [108] |
| Male | 75 | serutan® (psyllium granules) | No | Insufficient water intake after bulk-producer ingestion | [109] |
| Male | 81 | Psyllium-based Bulk laxative "Perdiem" | No | Insufficient water intake after bulk-producer ingestion | [110] |
| Male | 72 | Perdiem granules | No | Insufficient water intake after bulk-producer ingestion | [111] |
| Woman | 67 | Perdiem granules | No | Insufficient water intake after bulk-producer ingestion | [111] |
| Woman | 41 | Perdiem granules | No | Insufficient water intake after bulk-producer ingestion | [111] |
| Male | 66 | Plantago ovata seed (psyllium seed) | No | Insufficient water intake after bulk-producer ingestion | [112] |
| Male | 56 | Vegan diet, most probably chia seeds bezoar. | No | Strict vegan diet 15 Kg weight loss in 12 months. | [113] |
| Infant | 1.5 | Psyllium, beet & lentils | | Down syndrome | [146] |

| Gender | Age (years) | phytobezoar Type | previous surgery | Other risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Man | 50 | Konjac bezoar | | | [141] |
| Man | 60 | Mushroom | | | [149] |

**Table 5: list for unusual food, pulpy and non-fibrous food bezoar cases**

| Gender | Age | phytobezoar Type | previous surgery | Risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Male | 50 Y | Male, Fatty acids and Lecithin | No | No | [114] |
| Female | 44 Y. | large amounts of honeycomb | No | Irritable bowel syndrome Consumption of large quantities of honeycomb for health benefits | [115] |
| Male | 65 Y | Chestnuts phytobezoar | No | Chestnuts consumption | [116] |
| Male | 61 Y. | Chestnuts phytobezoar | No | Consumption of large quantity of Chestnuts | [117] |
| Male | 1.3 Y (16 months) | dried Apricots | No | whole dried apricots , inadequate mastication | [118] |
| Male | 78 Y. | phytobezoar (undigested Apple pieces & whole maize grains) | No | abnormal mastication habits as well as a jejunal diverticulum | [119] |
| Male | 86 | pickled plume | No | accidently ingested japanese pickled plum with seeds. | [120] |
| Male | 69 | dried Apple | No | coronary arteries bypass, diabetes and hypertension. He had no teeth, and not used dentures for the past 10 years. Only chew various foods with his gums only. but admitted to swallowing food in whole occasionally. | [121] |
| Not specified | Not specified | 150 cases, GI obstruction due to different types of phytobezoars (orange pulp, figs, apples, grapefruit, persimmons, coconuts, cabbage, beans and potatoes) | All have had previous gastrectomy for duodenal ulcer. | | [9] |
| Female | 62 Y. | phytobezoar | No | many years she had eating quite a substantial proportion of raw or poorly cooked long stemmed vegetables. | [122] |
| Male | 4 Y. | Undigested rhubarb | no | undigested rhubarb | [78] |

| Gender | Age | phytobezoar Type | previous surgery | Risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Male | 75 | Meat bolus bezoar | No | Our patient had not gone under gastric surgery but recalled a right inguinal hernia repair40~ years ago, had a history of diabetes mellitus type II, and he wears a dental prosthesis which increases the possibility of bezoar formation in his GIT. | [123] |
| Female | 59 | swollwed whole potatoes | No | medical history of cerebral palsy due to birth hypoxia | [124] |
| Female | 53 | swallowed whole potatoes | No | Medical history of learning disability, autism and partial sightedness, | [124] |
| 2 Males, 1 Female | 0.25 - 0.3 Y (3-4 months)2 girl, 1 boy | potato bezoar | No | mashed potato | [125] |
| 5 Male, 9 Female | 52 -82 Y | Rice cake bezoar , 4 cases no surgery | (7 cases , 3 Male & 4 females) 4 appendictomy (2F+2M), 1 hysterctomy, (1, M) gasterectomy, 1 ileus. +3 cases female cesarian resections | All patients had ingested rice cake by swallowing without chewing. | [37] |
| 5 Male, 3 Female | 62, 63, 68, 76, 85 (M); 56, 65, 83 (F) | Rice cake bezoar, 6 cases no surgery | Yes, 2 cases (1 m and 1F*) abdominal syrgeries | 5 have dentures (3M+2F), 1 quick feeder, 1had operation, 1 operation and quick feeder (F*) | [39] |
| Female | 68 Y | Rice cake bezoar | No | She had eaten a toasted rice cake without chewing well because of her dentures the day before. | [126] |
| Female | 59 Y | Rice cake bezoar | No | She had eaten a toasted rice cake without chewing well because of her dentures the day before. | [126] |
| Female | 70 Y | Rice cake bezoar | No | She had dentures and had eaten a boiled rice cake 5 hours previously. | [126] |
| Male | 37 | Rice cake bezoar | No | He had eaten a boiled rice cake without chewing well the day before. | [126] |

| Gender | Age | phytobezoar Type | previous surgery | Risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Male | 64 | Potato, brussels sprouts, and broccoli | No | for obesity and prior cholecystectomy. | [127] |
| 9 Males/ 8 Females | (44-80) | Rice cake bezoar, 10 cases with no surgery | Appendicectomy (3 cases), Hysterectomy (4 cases) | Gulping habit (2 cases), toothless (1case), | [54] |

**Table 6: list of phytobezoar cases due to chronic diseases or GI disorders**

| Gender | Age | phytobezoar Type | previous surgery | Risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| 1 Boy, 5 Girls | (12-18 Y) | Phytobezoar | | Dysautonomia (6 patients) | [128] |
| Male | 60 | food bezoar | No | GI laziness and constipation due to khat consumption | [129] |
| Female | 24 | Giant phytobezoar | No | No expected risk factors., but most likely dyspepsia and hypoacidity (she mentioned consumption of antacids for three months before admission). These factors may causes delayed gastric emptying. | [130] |
| Housewife | 41 | phytobezoar formed of undigested vegetable matter (peas), fibrous material and decomposed food residue. | No | No (may be poor physical exercise leads to GI laziness) | [91] |
| Male | 11 | Boy, 11 Y., phytobezoar | No | Intellectual disability and seizure disorder | [131] |
| Female | 21 | phytobezoar (Food matter) | | Bulimia nervosa Binge eating episodes | [74] |
| Male | 48 | a huge vegetable materials | No | hypertension and kidney failure , high fiber diet | [86] |
| Female | 19 | (Vegetable matter) | No | Anorexia nervosa , Binge-purge , Hematemesis | [40] |
| Girl | 11 | giant phytobezoar due to ileal duplication. | No | Ileal duplication. | [132] |
| Woman | 59 | phytobezoar | No | type 2 diabetes and hypertension | [133] |
| (20 males, 24 females) | (55.36±15.17) | | 7 cases out of 44 patients, have had previous surgeries | Chronic diseases (13 cases) out of 44 patients | [34] |
| 90 Females & 63 M ales including (29 girls + 54 Boys) | Age range (2-80 Y.). | | Only 10 surgery cases | gastrointestinal strictures (10 cases/153, 6.5%), diabetes (4 cases/153, 2.6%) and neuropsychiatric disease (4 cases/153, 2.6%). | [35] |

| Gender | Age | phytobezoar Type | previous surgery | Risk factors & medical history | Ref. |
|---|---|---|---|---|---|
| Man | 57-years | phytobezoar | No | type 2 diabetes with autonomic neuropathy have impaired gastrointestinal emptying functions, increasing the retention time of food and causing phytobezoar | [134] |
| Man | 84-years | phytobezoar | No | Ischemic HD, Coronary artery by-pass graft, heart failure, Dementia, diabetes, benign prostatic hypertrophy, bipolar disorder. | [140] |
| Girl | 17-Years | phytobezoar | No | Chronic eating disorder | [136] |
| Male | 67-Year | Phytobezoar | Yes | Surgery due to Prostatic cancer - radiotherapy | [147] |
| Boy | 12-Years | phytobezoar | No | Meckel's diverticulum | [150] |

## 7. References


1. *Abu-Jkeim, N.; Al Hazmi, A.; Ibrahim, R.; Abu Damis, A.; Tawfik, S.; Mansour, M.; Alawad, A.M. A. Small bowel obstruction due to phytobezoar and adhesions following laparoscopic cholecystectomy. International Journal of Medicine. 2015, 3(2), 63-65 https://doi.org/10.14419/ijm.v3i2.4943*
2. *Akin, I. B., Altay, C., Egeli, T., Yilmaz, I., & Obuz, F. B. (2020). Rare Cause of Small Bowel Obstruction: Multiple Phytobezoars. Cyprus Journal of Medical Sciences, 5(3), 277-279. https://doi.org/10.5152/cjms.2020.1164*
3. *Arsalan, O.; Hussein, A.; El Malik, S. A case of antroduodenal phytobezoar following laparoscopic sleeve gastrectomy. Hamdan Med. J. 2014, 7, 117–120. https://doi.org/10.7707/hmj.v7i1.305.*
4. *Aryannezhad, S.; Sadeghian, Y.; Shapoori, P.; Valizadeh, M.; Barzin, M. A rare case report of late-onset phytobezoar formation following laparoscopic sleeve gastrectomy: Delayed redo bariatric surgery. BMC Surg. 2021, 21, 254. https://doi.org/10.1186/s12893-021-01254-8*
5. *Ashfaq, A.; Madura, J.A. 2nd; Chapital, A.B. An unusual case of biliary bezoar causing small bowel obstruction in a patient with ampullary diverticulum and stapled gastroplasty. BMJ Case Rep. 2014, bcr2014207455. https://doi.org/10.1136/bcr-2014-207455.*
6. *Ben-Porat, T.; Elazary, R.; Goldenshluger, A.; Sherf-Dagan, S.; Grinbaum, R.; Beglaibter, N. Phytobezoar: A rare late complication following laparoscopic sleeve gastrectomy surgery. Int. Surg. J. 2017, 4, 1803–1805. https://doi.org/10.18203/2349-2902.isj20171644.*
7. *Ben-Porat, T.; Sherf-Dagan, S.; Goldenshluger, A.; Yuval, J.B.; Elazary, R. Gastrointestinal phytobezoar following bariatric surgery: Systematic review. Surg Obes Relat Dis. 2016, 12(9), 1747-1754. https://doi.org/10.1016/j.soard.2016.09.003.*
8. *Deront Bourdin, F.; Iannelli, A.; Delotte, J. Phytobezoar: an unexpected cause of bowel obstruction in a pregnant woman with a history of Roux-en-Y gastric bypass. Surg Obes Relat Dis. 2014, 10(6), e49-51. https://doi.org/10.1016/j.soard.2014.07.009.*
9. *Buchholz, R.R.; Haisten, A.S. Phytobezoars following gastric surgery for duodenal ulcer. Surg Clin North Am. 1972, 52(2), 341-52. https://doi.org/10.1016/s0039-6109(16)39686-4.*
10. *Cabo-Navarro, D.; Fort López-Barajas, J.M.; Mesa Manteca, J.; Lecube Torelló, A. Fitobezoar en remanente gástrico tras cirugía bariátrica [Fitobezoar in gastric pouch after bariatric surgery]. Med Clin (Barc). 2009, 132(5), 200. Spanish. https://doi.org/10.1016/j.medcli.2008.09.008.*
11. *Chisholm, E.M.; Leong, H.T.; Chung, S.C.; Li, A.K. Phytobezoar: an uncommon cause of small bowel obstruction. Ann R Coll Surg Engl. 1992, 74(5), 342-4.*
12. *Claro, M.; Santos, D. C.; Silva, A. A.; Deus, C.; Grilo, J.; Sousa, D.; Martins, J. A. (). When eating makes you sick–Gastric stump obstruction caused by a phytobezoar. A case report and literature review. International Journal of Surgery Case Reports. 2021, 79, 263-266. https://doi.org/10.1016/j.ijscr.2021.01.034*
13. *Cooper, S.G.; Tracey, E.J. Small-bowel obstruction caused by oat-bran bezoar. N Engl J Med. 1989, 320 (17), 1148-9. https://doi.org/10.1056/NEJM198904273201717.*
14. *Cortés, C.; Silva, C.;, Gastric bezoar as complication of gastric banding. Report of one case. Revista Medica de Chile. 2008, 136, (11),*

*1457–1459. https://doi.org/10.4067/s0034-98872008001100013.*

15. *Das, S.S.; AbdelAziz, Z.; Bondok, W.Z.A.M.; Juma, F.I.B.; Khatib, F.H.A. A Phytobezoar Causing Terminal Ileal Obstruction Following Revision Bariatric Surgery: A Case Report. Cureus. 2023, 15(4), e37353. https://doi.org/10.7759/cureus.37353.*
16. *de Toledo, A.P.; Rodrigues, F.H.; Rodrigues, M.R.; Sato, D.T.; Nonose, R.; Nascimento, E.F.; Martinez, C.A. Diospyrobezoar as a cause of small bowel obstruction. Case Rep Gastroenterol. 2012, 6(3), 596-603. https://doi.org/10.1159/000343161.*
17. *Dray, X.; Samaha, E.; Tuszynski, T. Pouch outlet obstruction by psyllium-based bulk laxative bezoar in a patient with gastric banding. Surg Obes Relat Dis. 2015, 11(2), 483-4. https://doi.org/10.1016/j.soard.2015.01.018.*
18. *Edelstein, M.M.; Freed, E.; Wexler, M. Diospyrobezoar of the jejunum in a postgastrectomy patient. Arch Surg. 1971, 103(6):765-6. https://doi.org/10.1001/archsurg.1971.01350120129026.*
19. *Ertugrul, I.; Tardum-Tardu, A.; Tolan, K.; Kayaalp, C.; Karagul, S.; Kirmizi, S. Gastric bezoar after Roux-en-Y gastric bypass for morbid obesity: A case report. Int J Surg Case Rep. 2016, 23:112-5. https://doi.org/10.1016/j.ijscr.2016.04.008.*
20. *Erzurumlu K, Malazgirt Z, Bektas A, Dervisoglu A, Polat C, Senyurek G, Yetim I, Ozkan K. Gastrointestinal bezoars: a retrospective analysis of 34 cases. World J Gastroenterol. 2005 Mar 28;11(12):1813-7. https://doi.org/10.3748/wjg.v11.i12.1813.*
21. *Gachabayov, M.; Abdullaev, A.; Mityushin, P.; Gilyazov,T. "Each worm to his taste: some prefer to eat nettles – a giant gastric phytobezoar," Clinical Case Reports. 2016, 4, 710-711. https://doi.org/10.1002/ccr3.593*
22. *Geerts, B.; Verstraeten, T.; Vanslambrouck, K.; Vandenhaute, S.; Dillemans, B.; Coenegrachts, K. Unusual Cause of Small Bowel Obstruction Caused by a Prune Pit. J Belg Soc Radiol. 2015, 99(1), 95-96. https://doi.org/10.5334/jbr-btr.838.*
23. *Gero, D.; Chosidow, D.; Marmuse, J.P. An unusual cause of solid food intolerance and gastric pouch dilation 5 years after Roux-en-Y gastric bypass: agar-agar bezoar. Surg Obes Relat Dis. 2015, 11(5), 1184-5. https://doi.org/10.1016/j.soard.2015.05.010.*
24. *Goitein, D.; Papasavas, P.K.; Gagné, D.J.; Caushaj, P.F. Late perforation of the jejuno-jejunal anastomosis after laparoscopic Roux-en-Y gastric bypass. Obes Surg. 2005, 15(6), 880-2. https://doi.org/10.1381/0960892054222821.*
25. *Grande, G.; Manno, M.; Zulli, C.; Barbera, C.; Mangiafico, S.; Alberghina, N.; Conigliaro, R.L. An alternative endoscopic treatment for massive gastric bezoars: Ho:YAG laser fragmentation. Endoscopy. 2016, 48 Suppl 1, E217. https://doi.org/10.1055/s-0042-109057.*
26. *Hai, A.T.T.; Thi, L.N.K.; Tuyet, C.P.T.; Ngoc, Y.M.; Vu, O.H.;Thi, O.N.; Thi, H.H., Thuy, L.N. Nutritional Status and Diet Characteristics in Patients with Intestinal Obstruction Due to Food Residues at Hospital in Vietnam. Asian J Dietetics. 2022, 19, 9. https://jnl.calorie-smile.jp/eng/wordpress/wp-content/uploads/2022/04/asian_journal_of_dietetics_4_1_2022_9_13.pdf*
27. *Hemmasi, G.; Zanganeh, E.; Hosseini, S.A.; Alimadadi, M.; Ahmadi, A.; Hajiebrahimi, S.; Seyyedmajidi, M. Risk factors, endoscopic findings, and treatments in upper gastrointestinal bezoars: multi-center experience in Iran. Gastroenterol Hepatol Bed Bench. 2021, 14(2), 160-164.*
28. *Ionescu, A.M.; Rogers, A.M.; Pauli, E.M.; Shope, T. R. (). An unusual suspect: Coconut bezoar after laparoscopic Roux-en-Y gastric bypass. Obesity Surgery, 2008,18(6), 756-758. https://doi.org/10.1007/s11695-007-9343-4*
29. *Islam, S.; Saroop, S.; Bheem, V.; Harnarayan, P.; Naraynsingh, V. Mango seed causing acute large bowel obstruction in descending colon-world's first reported case. Int J Surg Case Rep. 2018, 51, 125-129. https://doi.org/10.1016/j.ijscr.2018.08.028.*
30. *Jang, J.O.; Choi, C.W.; Ryu, D.G.; Park, S.B.; Kim, S. J. (). Clinical Outcomes of Patients With Gastric Phytobezoars: Implication of Drinking Carbonated Beverages as a Treatment. Korean J Helicobacter & Upper Gastrointestinal Research, 2024, 24(3), 259-266. https://doi.org/10.7704/kjhugr.2024.0032*
31. *Kosmidis, C.S.; Mystakidou, C.M.; Varsamis, N.; Koulouris, C.; Sevva, C.; Papadopoulou, K.; Michael, C.; Katsios, N.I.; Theodorou, V.; Miltiadous, P.; Papadopoulos, K.; Vlassopoulos, K.; Zarampouka, K.; Mantalovas, S.; Phytobezoar-Induced Mechanical Ileus and Incipient Intussusception: A Case Report. Medicina (Kaunas). 2023, 59(7),1227. https://doi.org/10.3390/medicina59071227.*
32. *Krausz, M.M.; Moriel, E.Z.; Ayalon, A.; Pode, D.; Durst, A.L. Surgical aspects of gastrointestinal*

*persimmon phytobezoar treatment. Am J Surg. 1986, 152(5), 526-30. https://doi.org/10.1016/0002-9610(86)90221-7.*

33. *Laird Birmingham, C.; Cardew, S.; Gritzner, S. "Gastric bezoar in anorexia nervosa," Eating and Weight Disorders. 2007, 12, e28–e29. https://doi.org/10.1007/BF03327586.*
34. *Liu, B; Kuang, S. Gastric Bezoars: A Retrospective Analysis of 44 Cases. Med Sci Monit. 2024, 30, e945377. https://doi.org/10.12659/MSM.945377.*
35. *Manatakis, D.K.; Acheimastos, V.; Antonopoulou, M.I.; Balalis, D.; Korkolis, D.P. Gastrointestinal Seed Bezoars: A Systematic Review of Case Reports and Case Series. Cureus. 2019, 11(5), e4686. https://doi.org/10.7759/cureus.4686.*
36. *Martines, G.; Musa, N.; Aquilino, F.; Picciariello, A.; Altomare, D.F. Sleeve gastrectomy combined with Nissen fundoplication as a single surgical procedure, is it really safe? A case report. Am. J. Case Rep. 2020, 21, e923543. https://doi.org/10.12659/AJCR.923543*
37. *Miura, T.; Kimura, N.; Nakamura, J.; Yamada, S.; Miura, T.; Yanagi, M.; Ikeda, Y.; Takaki, S.; Tani, Y.; Ito, T.; Nishihara, M.; Takahashi, T. Rice cake ileus--a rare and ethnic but important disease status in east-southern Asia. Intern Med. 2011, 50(22), 2737-9. https://doi.org/10.2169/internalmedicine.50.6152.*
38. *Ohya, Y.; Hayashida, S.; Yoneda, A.; Tsuji, A.; Inoue, T.; Chiyonaga, S.; Kuramoto, K.; Oda, K.; Iizaka, M.; Nakahara, O.; Inomata, Y. Laparoscopic-assisted treatment for diospyrobezoar-induced intestinal obstruction after distal gastrectomy and cholecystectomy. Drug Discov Ther. 2023, 17(3), 217-219. https://doi.org/10.5582/ddt.2023.01009.*
39. *Oka, A.; Amano, Y.; Uchida, Y.; Kagawa, K.; Takatori, K.; Kitajima, N.; Sonoyama, H.; Tada, Y.; Kusunoki, R.; Fukuba, N.; Oshima, N.; Moriyama, I.; Yuki, T.; Kawashima, K.; Ishihara, S.; Kinoshita, Y. [Small bowel obstruction and gastric ulceration resulting from rice cake ingestion -computed tomography diagnosis in eight patients-]. Nihon Shokakibyo Gakkai Zasshi. 2013, 110(10),1804-13.*
40. *Parameswaran, R.; Ferrando, J.; Sigurdsson, A. Gastric bezoar complicating laparoscopic adjustable gastric banding with band slippage. Obes Surg. 2006, 16(12),1683-4. https://doi.org/10.1381/096089206779319374.*
41. *Parsi, S.; Rivera, C.; Vargas, J.; Silberstein, M.W. Laparoscopic-assisted extirpation of a phytobezoar causing small bowel obstruction after Roux-en-Y laparoscopic gastric bypass. Am Surg. 2013, 79(2), E93-5.*
42. *Paschos, K.A.; Chatzigeorgiadis, A. Pathophysiological and clinical aspects of the diagnosis and treatment of bezoars. Ann Gastroenterol. 2019, 32(3), 224-232. https://doi.org/10.20524/aog.2019.0370.*
43. *Pinto, D.; Carrodeguas, L.; Soto, F.; Lascano, C.; Cho, M.; Szomstein, S.; Rosenthal, R. Gastric bezoar after laparoscopic Roux-en-Y gastric bypass. Obes Surg. 2006, 16(3), 365-8. https://doi.org/10.1381/096089206776116561.*
44. *Powell, S.D.; Pereira, N. Bezoars: Culprits of gastrointestinal obstruction that may lead to surgical intervention and potentially surgical complications. Clin Case Rep. 2023, 11(12), e8126. https://doi.org/10.1002/ccr3.8126.*
45. *Powers, W.F. 4th.; Miles, D.R. Phytobezoar causing small bowel obstruction seven years after laparoscopic Roux-en-Y gastric bypass. Surg Obes Relat Dis. 2011 Jul-Aug;7(4):e3-5. https://doi.org/10.1016/j.soard.2010.07.008.*
46. *Pratt, J.S.; Van Noord, M.; Christison-Lagay, E. The tethered bezoar as a delayed complication of laparoscopic Roux-en-Y gastric bypass: a case report. J Gastrointest Surg. 2007, 11(5), 690-2. https://doi.org/10.1007/s11605-007-0098-y.*
47. *Pujar,K.A.;Pai,A.S.;Hiremath,V.B.Phytobezoar: a rare cause of small bowel obstruction. J ClinDiagn Res. 2013, 7, 2298-299. http://dx.doi.org/10.7860/jcdr/2013/7248.3504*
48. *Qin, B.; Wan, X.L.; Guo, X.Y.; Dong, L. Successful endoscopic treatment of an intestinal diospyrobezoar migrated from the stomach. BMJ Case Rep. 2014, 2014, bcr2014204966. https://doi.org/10.1136/bcr-2014-204966.*

49. *Roy, M.; Fendrich, I.; Li, J.; Szomstein, S.; Rosenthal, R.J. Treatment option in patient presenting with small bowel obstruction from phytobezoar at the jejunojejunal anastomosis after Roux-en-Y gastric bypass. Surg Laparosc Endosc Percutan Tech. 2012, 22(4), e243-5. https://doi.org/10.1097/SLE.0b013e31825d6c07.*
50. *Sammut, S.J.; Majid, S.; Shoab, S. Phytobezoar: a rare cause of late upper gastrointestinal perforation following gastric bypass surgery. Ann R Coll Surg Engl. 2012, 94(2), e85-7. https://doi.org/10.1308/003588412X13171221588938.*
51. *Sarhan, M; Shyamali, B., Fakulujo, A.; Ahmed, L. Jejunal Bezoar causing obstruction after laparoscopic Roux-en-Y gastric bypass. J Soc Laparoendoscopic Surg. 2010, 14(4), 592-5. https://doi.org/10.4293/108680810X12924466008682.*
52. *Sealey, A.; J, Lau, N. S. Peanut phytobezoar: an unusual cause for small bowel obstruction. Journal of Surgical Case Reports. 2024, 8, 2024, rjae564. https://doi.org/10.1093/jscr/rjae564*
53. *Shiotsuki, K.; Imai, K.; Hotta, K. Small bowel obstruction by massive impacted diospyrobezoars relieved by endoscopic lithotripsy with a polypectomy snare using double balloon endoscopy. Dig Endosc. 2019, 31(6), e111-e112. https://doi.org/10.1111/den.13490.*
54. *Sugimoto, S.; Shimono, T.; Takeshita, T.; Yamamoto, A.; Shindo, D.; Miki, Y. Clinical and CT findings of small bowel obstruction caused by rice cakes in comparison with bezoars. Jpn J Radiol. 2019, 37(4), 301-307. https://doi.org/10.1007/s11604-019-00811-9.*
55. *Tabesh, E.; Dehghan, A.; Tahmasebi, M.; Javadi, N. Gastric phytobezoars as a very unusual cause of gastric outlet obstruction. J Res Med Sci. 2021, 26, 25. https://doi.org/10.4103/jrms.JRMS_115_20. .*
56. *Ueno, T.; Hanabata, N.; Katagai, A.; Okudera, R.; Arai, A.; Tomiyama, M. Phytobezoar Associated with Levodopa-carbidopa Intestinal Gel Infusion in Patients with Parkinson's Disease: A Case Report and Literature Review. Intern Med. 2021, 60(20), 3317-3320. https://doi.org/10.2169/internalmedicine.7210-21.*
57. *Vellar DJ, Vellar ID, Pucius R, Steedman PK. Phytobezoars--an overlooked cause of small bowel obstruction following vagotomy and drainage operations for duodenal ulcer. Aust N Z J Surg. 1986 Aug;56(8):635-8. https://doi.org/10.1111/j.1445-2197.1986.tb04519.x.*
58. *Verma, H.; Burns, D.; Nepomnayshy, D. Coconut Bezoars in Patients with Prior Bariatric Surgery: Report of Two Cases. Am J Gastroenterol. 2013, 108 (1), S220.*
59. *White, N.B.; Gibbs, K.E.; Goodwin, A.; Teixeira, J. Gastric bezoar complicating laparoscopic adjustable gastric banding, and review of literature. Obes Surg. 2003, 13(6), 948-50. https://doi.org/10.1381/096089203322618849*
60. *Yang, S. ; Cho, M. J. Clinical characteristics and treatment outcomes among patients with gastrointestinal phytobezoars: a single-institution retrospective cohort study in Korea. Frontiers in Surgery. 2021, 8, 691860 https://doi.org/10.3389/fsurg.2021.691860*
61. *Zapata, R.; Castillo, F.; Córdova, A. Bezoar gástrico como complicación de la cirugía de obesidad mórbida. Caso clínico y revisión de la bibliografía [Gastric food bezoar as a complication of bariatric surgery. Case report and review of the literature]. Gastroenterol Hepatol. 2006, 29(2), 77-80. https://doi.org/10.1157/13083903.*
62. *Conomea, J.R.; Bobak, A.K.; Kambhampati, H.; Stauffer, J.; Herman, M. Duodenal Phytobezoar Treated With Endoscopic Removal: A Case Report. Cureus. 2025, 17(2), e79147. https://doi.org/10.7759/cureus.79147.*
63. *Patiño-Araujo, B.; Salgado-Báez, M.E.; Villavicencio-Logroño, G.P.; Ayora-Pérez, M.J.; Zapata-Jaramillo, G.S.; Salgado-Macías, N. Early small bowel obstruction by phytobezoar related to zopiclone induced amnesia after revisional Roux-en-Y gastric bypass: a case report. Journal of Surgical Case Reports. 2025, 4, rjaf202. https://doi.org/10.1093/jscr/rjaf202*
64. *DeBakey, M.; Oschner, A. Bezoars and concretions. Surgery. 1939, 5, 132-160.*
65. *Dolan, P.A.; Thompson, B.W. Management of persimmon bezoars (diospyrobezoars). South Med J. 1979, 72(12), 1527-8, 1531. https://doi.org/10.1097/00007611-197912000-00010.*
66. *Ha, S.S.; Lee, H.S.; Jung, M.K.; Jeon, S.W.; Cho, C.M.; Kim, S.K.; Choi, Y.H. Acute intestinal obstruction caused by a persimmon phytobezoar after dissolution therapy with Coca-Cola. Korean J Intern Med. 2007, 22(4), 300-3. https://doi.org/10.3904/kjim.2007.22.4.300.*
67. *Odeh, A.M.; Alkhalifa, A.A.; AlHajji, M.A.; Alahmed, A.J.; Alsalman, J.H.; AlMulhem, A.M.; AlGhadeer, M.S.; Kurdi, E.; Albader, F.S.; Alsuwaigh, A.; Khan, M.A. A Rare Case of Gastric Phytobezoar (Diospyrobezoar) in a Healthy Adult. Cureus. 2024, 16(8), e68353. https://doi.org/10.7759/cureus.68353.*
68. *O'leary, C.M. Diospyrobezoar: A review of fourteen cases with an analysis of forty-six collected cases from the literature. AMA Arch Surg. 1953, 66(6), 857–868. https://doi.*

*org/10.1001/archsurg.1953.01260030877016.*

69. *Zheng, Y.X.; Prasoon, P.; Chen, Y.; Hu, L.; Chen, L. ''Sandwich" treatment for diospyrobezoar intestinal obstruction: a case report. World J Gastroenterol. 2014, 20(48), 18503-6. https://doi.org/10.3748/wjg.v20.i48.18503.*
70. *Alexandre, A.R.; Costa, L.D.; Raimundo, P. Giant granadilla's seeds phytobezoar rectal impactation: a very unusual case of intestinal obstruction. BMJ Case Rep. 2018, 2018, bcr2017223176. https://doi.org/10.1136/bcr-2017-223176.*
71. *Ali, A.Y.; Biyikli, A.; Mohamed, A.A.; Dhaley, A.; İbrahim, İ.G. Infrequent cause of small bowel obstruction in a child: A lemon. Ann Med Surg (Lond). 2022, 76, 103488. https://doi.org/10.1016/j.amsu.2022.103488.*
72. *Cindy, M.; Blanco, O.; Mariela, L. P.; Berger, M.; Miryam, M. Bezoar in Pediatrics: Case Series and Literature Review. Science Volks Open Access Paediatrics. 2024, 3(5), 166-170. https://doi.org/10.58624/SVOAPD.2024.03.083*
73. *Eitan, A.; Bickel, A.; Katz, I.M. Fecal impaction in adults: report of 30 cases of seed bezoars in the rectum. Dis Colon Rectum. 2006, 49(11), 1768-71. https://doi.org/10.1007/s10350-006-0713-0.*
74. *Fazio, R. M.; Shah, P.; Soe, E.; Iswara, K.; Chen, I. "Gastric bezoar causing massive gastric distention and functional outlet obstruction in a patient with bulimia nervosa," Journal of Medical Cases. 2016, 7, 7, 312–314. https://doi.org/10.14740/jmc2560w*
75. *FDA, 2007: Food and Drug Administration, HHS. Laxative drug products for over-the-counter human use; psyllium ingredients in granular dosage forms. Final rule. Fed Regist. 2007, 72(60), 14669-74. https://www.govinfo.gov/content/pkg/FR-2007-03-29/pdf/E7-5740.pdf*
76. *Fernando, A.R.; Bulathsinghela, R.; Samarasekera, D.N. An unusual cause of small bowel obstruction due to an ingested mango seed: a case report. BMC Res Notes. 2017, 10(1):549. https://doi.org/10.1186/s13104-017-2875-3.*
77. *Gentile, M.; Vergara, L.; Schiavone, V.; Cestaro, G.; Sivero, L. Harry Potter's Occlusion: Report of a Case of Pumpkin Seed Bezoar Rectal Impact. Front Surg. 2022, 9, 902701. https://doi.org/10.3389/fsurg.2022.902701.*
78. *Glatstein, M.; Danino, D.; Rimon, A.; Keidar, S.; Scolnik, D. An unusual cause of small bowel obstruction in a child: ingested rhubarb. Case Rep Surg. 2013, 2013, 497214. https://doi.org/10.1155/2013/497214.*
79. *Manatakis, D.; Sioula, M.; Passas, I.; Zerbinis, H.; Dervenis, C. Rectal seed bezoar due to sunflower seed: a case report and review of the literature. Pan Afr Med J. 2018, 31,157. https://doi.org/10.11604/pamj.2018.31.157.12539.*
80. *Moons, P.; Dikken, F.P.; Raven, E.E.; Tan, K.G. Ernstige obstipatie door het eten van ongeschilde zonnebloempitten [Severe obstipation due to eating unshelled sunflower seeds]. Ned Tijdschr Geneeskd. 2000, 144(39), 1878-9. Dutch.*
81. *Purcell, L.; Gremse, D.A. Sunflower seed bezoar leading to fecal impaction. South Med J. 1995 Jan;88(1):87-8. https://doi.org/10.1097/00007611-199501000-00013.*
82. *Samuel, S.; Sharko, A.; Shrestha, J.; Sherchan, R.; Baig, S. A Rare Case of Small Bowel Obstruction Due to Sunflower Seeds Consumption in an Adult. Cureus. 2021, 13(4), e14330. https://doi.org/10.7759/cureus.14330.*
83. *Thing, B.A.; Jørgensen, H. Rektal bezoar forårsaget af solsikkekerner [Rectal bezoar caused by sunflower seeds]. Ugeskr Laeger. 2010, 172(42), 2905-6.*
84. *Salada, K.O.; Jacobson, E. Seeds of Discomfort: An Unusual Case of Pediatric Abdominal Pain. Cureus. 2023, 15(12), e50625. https://doi.org/10.7759/cureus.50625.*
85. *Marchese, S.; Bertucci, B.; Manti, F.; Berritto, D.; Roperto, A.g.; Tamburrini, S. Rectal impaction due to prickly pear seeds bezoar: a case report. J biol regul homeost agents. 201, 29(3), 707-11.*
86. *Hassranah, S.C.; Ramnarine, A.; Parbhu, S.; Naraynsingh, V. Small Bowel Obstruction Caused by an Aggressive Weight Loss Diet in a Patient With No Predisposing Factors. Cureus. 2022, 14(12):e32594. https://doi.org/10.7759/cureus.32594.*
87. *Hefny, A.F.; Ayad, A.Z.; Matev, N.; Bashir, M.O. Intestinal obstruction caused by a laxative drug (Psyllium): A case report and review of the literature. Int J Surg Case Rep. 2018, 52, 59-62. https://doi.org/10.1016/j.ijscr.2018.10.001.*
88. *Kavlakoğlu, B.; Oral, S. Very rare coincidence: perforated duodenal ulcer and olive seed phytobezoar. Turk J Gastroenterol. 2011, 22(4):442-3. https://doi.org/10.4318/tjg.2011.0294.*
89. *Khan, S.; Jiang, K.; Zhu, L.P.; Khan, I.A.; Ullah, K.; Khan, S.; Chen, X.; Wang, B.M. Upper Gastrointestinal Manifestation of Bezoars and the Etiological Factors: A Literature Review. Gastroenterol Res Pract. 2019, 5698532. https://doi.org/10.1155/2019/5698532.*

90. *Shaw, A.G.; Peacock, O.; Lund, J.N.; Tierney, G.M.; Larvin, M.; Speake, W. Large bowel obstruction due to sesame seed bezoar: a case report. J Med Case Rep. 2007, 1, 159. https://doi.org/10.1186/1752-1947-1-159.*
91. *Tiwari, A; Gupta, V.; Hazrah, P.; Chaudhary, A.K.; Lal, R.: Phy-tobezoar: a rare cause of acute small bowel obstruction in an innocent abdomen. Arq Bras Cir Dig. 2013, 26, 342-343. http://dx.doi.org/10.1590/S0102-67202013000400020.*
92. *Su, W.; Chen, J.; Zhao, Y.; Xu, P.; Yuan, J.; Luo, C.; Liu, J.; Wang, B. Diagnosis and treatment of giant colonic fecalith in a child: a case report. Front Pediatr. 2025, 13, 1598807. https://doi.org/10.3389/fped.2025.1598807.*
93. *Schoeffl, V.; Varatorn, R.; Blinnikov, O.; Vidamaly, V. Intestinal obstruction due to phytobezoars of banana seeds: a case report. Asian J Surg. 2004, 27(4), 348-51. https://doi.org/10.1016/s1015-9584(09)60067-1.*
94. *Yigzaw, K.T.; Mulualem, D.A.; Asfaw, B.A.; Moges, M.T.; Tegegne, B.A.; Shibesh, M.H.; Wassie, M.T. Uncommon cause of large bowel obstruction in a pediatric patient: A case of phytobezoar induced by peas. Radiol Case Rep. 2025, 20(5), 2393-2398. https://doi.org/10.1016/j.radcr.2025.01.097.*
95. *Wu, X.; Zhang, W.; Xia, M. Phytobezoar-induced intestinal obstruction relieved via enteroscopy. Rev Esp Enferm Dig. 2025, 10.17235/reed.2024.10976/2024. Advance online publication. https://doi.org/10.17235/reed.2024.10976/2024.*
96. *Ozcan, U.A.; Yilmaz, S.; Akansel, S.; Cizmeli, M.O.; Ertem, M. An unusual cause of small bowel obstruction: CT diagnosis of dried apricots. Emerg Radiol. 2007, 14(6):417-9. https://doi.org/10.1007/s10140-007-0625-x.*
97. *Atila, K.; Güler, S.; Bora, S.; Gülay, H. An unusual cause of small bowel perforation: apricot pit. Ulus Travma Acil Cerrahi Derg. 2011, 17(3), 286-8.*
98. *Al-Najjar, M.; Arthur, T. The nectarine pit as a cause for small bowel obstruction and perforation: a case report. Case Rep Surg. 2013, 2013, 902943. https://doi.org/10.1155/2013/902943.*
99. *McClurken, J.B.; Carp, N.Z. Bran-induced small-intestinal obstruction in a patient with no history of abdominal operation. Arch Surg. 1988, 123(1).; 98-100. https://doi.org/10.1001/archsurg.1988.01400250108021.*
100. *Rosario, P.G.; Gerst, P.H.; Prakash, K.; Albu, E. Dentureless distention: oat bran bezoars cause obstruction. J Am Geriatr Soc. 1990, 38(5), 608. https://doi.org/10.1111/j.1532-5415.1990.tb02419.x.*
101. *Rawl, R ; Browne, L. Watch It Grow: Esophageal Impaction With Chia Seeds: 833. Am J of Gastroenterol. 2014, 109(1), S244-S245,*
102. *Salguero Molpeceres, O.; Seijas Ruiz-Coello, M.C.; Hernández Núñez, J.; Caballos Villar, D.; Díaz Picazo, L.; Ayerbe García-Monzón, L. Obstrucción esofágica por la fibra dietética Plantago ovata, una complicación prevenible mediante la información [Esophageal obstruction caused by dietary fiber from Plantago ovata, a complication preventable by adequate information]. Gastroenterol Hepatol. 2003, 26(4), 248-50.*
103. *Schlag, S.; Liese, J.; Veldhoen, S. Akutes Abdomen mit galligem Erbrechen – Ein Leinsamen-Bezoar als ungewöhnliche Ursache eines obstruktiven Ileus bei einer Adoleszenten mit Anorexie [Acute abdomen with Bilious Vomiting - A Linseed Bezoar as a Rare Cause of an Obstructive Ileus in an Anorexic Adolescent]. Klin Padiatr. 2021, 233(2), 72-74. https://doi.org/10.1055/a-1170-7435.*
104. *Shin, S.; Kim, J.H.; Mun, Y.H.; Chung, H.S. Acute esophageal obstruction after ingestion of psyllium seed husk powder: A case report. World J Clin Cases. 2022, 10(7), 2336-2340. https://doi.org/10.12998/wjcc.v10.i7.2336.*
105. *Yeoh, S.W. Esophageal bezoar due to karaya gum granules used as a laxative. Clin J Gastroenterol . 2017,10, 437–441. https://doi.org/10.1007/s12328-017-0764-x*
106. *Fisher, R.E. Psyllium Seeds: Intestinal Obstruction. Cal West Med. 1938, 48(3),190.*
107. *Herrle, F.; Peters, T.; Lang, C.; von Fluee, M.; Kern, B.; Peterli, R. Bolus obstruction of pouch outlet by a granular bulk laxative after gastric banding. Obes Surg. 2004, 14(7), 1022-4. https://doi.org/10.1381/0960892041719563.*
108. *Brown, D.C.; Doughty, J.C.; George, W.D. Surgical treatment of oesophageal obstruction after ingestion of a granular laxative. Postgrad Med J. 1999, 75(880), 106. https://doi.org/10.1136/pgmj.75.880.106.*
109. *Hinkel, C.L. Complete obstruction of the esophagus following serutan ingestion. J Am Med Assoc. 1951, 146(12), 1129-31. https://doi.org/10.1001/jama.1951.63670120005009b.*
110. *Noble, J.A.; Grannis, F.W. Acute esophageal obstruction by a psyllium-based bulk laxative. Chest. 1984, 86(5):800. https://doi.org/10.1378/chest.86.5.800a.*

111. *Schneider, R.P. Perdiem causes esophageal impaction and bezoars. South Med J. 1989, 82(11), 1449-50. https://doi.org/10.1097/00007611-198911000-00032.*

112. *Pérez-Piqueras, J.; Silva, C.; Jaqueti, J.; Sáez, M.A.; Martinez, D.; Santa, J.M.; Peralba, I. Endoscopic diagnosis and treatment of an esophageal bezoar resulting from bulk laxative ingestion. Endoscopy. 1994, 26(8), 710. https://doi.org/10.1055/s-2007-1009074.*

113. *Ho, D.; Nahm, C.B. Phytobezoar mimicking superior mesenteric artery syndrome in acute duodenal obstruction: a case report and review of literature. J Surg Case Rep. 2025, 2025(6), rjaf358. https://doi.org/10.1093/jscr/rjaf358.*

114. *Hsu, H. H.; Grove, W. E.; Mindulzun, R.; Knauer, C. M. "Gastric bezoar caused by lecithin: an unusual complication of health faddism," The American Journal of Gastroenterology, 1992, 87, 794–796. PMID: 1590324.*

115. *Katsinelos, P.; Pilpilidis, I.; Chatzimavroudis, G.; Katsinelos, T.; Lazaraki, G.; Fasoulas, K.; Zavos, C.; Kountouras, J. Huge gastric bezoar caused by honeycomb, an unusual complication of health faddism: a case report. Cases J. 2009, 2, 7077. https://doi.org/10.1186/1757-1626-2-7077.*

116. *Okagawa, Y.; Takada, K.; Arihara Y.; Kato, J. "A case of gastric perforation caused by chestnut bezoars," Nihon Shokakibyo Gakkai Zasshi= The Japanese Journal of Gastro-Enterology. 2017, 114, 1830–1835. https://doi.org/10.11405/nisshoshi.114.1830*

117. *Ravindra, R.K.; Das, A.; Chew, G.L.; Daniel, E. Small bowel obstruction with multiple perforations post chestnut ingestion. BMJ Case Rep. 2019, 12(2), bcr-2018-227999. https://doi.org/10.1136/bcr-2018-227999.*

118. *Piotto, L.; Gent, R. Dried apricots: an unusual cause of bowel obstruction. Pediatr Radiol. 2005 Dec;35(12):1224-6. https://doi.org/10.1007/s00247-005-1552-1.*

119. *Tayeb, M.; Khan, F.M.; Rauf, F.; Khan, M.M. Phytobezoar in a jejunal diverticulum as a cause of small bowel obstruction: a case report. J Med Case Rep. 2011, 5, 482. doi: 10.1186/1752-1947-5-482*

120. *Tsuchida, T.; Ishizuka, K.; Ohira, Y. Intestinal Obstruction Caused by Japanese Pickled Plum. J Gen Intern Med. 2023, 38(5), 1303. https://doi.org/10.1007/s11606-023-08046-1.*

121. *Ooi, S.; Hong, K. Small bowel obstruction caused by dried apple. Int J Surg Case Rep. 2015, 10, 154-7. https://doi.org/10.1016/j.ijscr.2015.03.038.*

122. *De Cesare, A.; Fiori, E.; Bononi, M.; Ferraro, D. Phytobezoar-induced small bowel obstruction associated with a concomitant gastric phytobezoar and ulcer in an elderly woman. Ann Ital Chir. 2015, 86(1), 70-7.*

123. *Albostani A, Kfelati F, Alsaadi W, Faraman RA, Farman A. Small bowel obstruction due to a meat bolus bezoar: the second case report in literature. Ann Med Surg (Lond). 2024;86(2):1139-1143. Published 2024 Jan 3. https://doi.org/10.1097/MS9.0000000000001633*

124. *Castren, E.; Hakeem, A.; Mahmood, N.S.; Aryal, K. Two cases of small bowel obstruction secondary to a swallowed potato. BMJ Case Rep. 2015, 2015, bcr2015213744. https://doi.org/10.1136/bcr-2015-213744.*

125. *Chao, H.C.; Chang, K.W.; Wang. C.J. Intestinal obstruction caused by potato bezoar in infancy: a report of three cases. Pediatr Neonatol. 2012, 53(2), 151-3. https://doi.org/10.1016/j.pedneo.2012.01.015.*

126. *Oka, A.; Ishihara, S.; Mikami, H.; Sonoyama, H.; Mishiro, T.; Tobita, H.; Kawashima, K.; Miyake, T.; Ishimura, N.; Furuta, K.; Kinoshita ,Y.; Nishina, M. Retained Rice Cake: A Unique Upper Gastrointestinal Foreign Body: Case Report and a Literature Review. Intern Med. 2019, 58(17), 2485-2494. https://doi.org/10.2169/internalmedicine.2760-19.*

127. *Serrano, K.D.; Tupesis, J.P. Small bowel obstruction from potato and broccoli phytobezoar mimicking mesenteric ischemia. J Emerg Med. 2013, 44(1), 79-81. https://doi.org/10.1016/j.jemermed.2011.06.018.*

128. *Shah, M.; Wan, D.Q.; Liu, Y.; Rhoads, J.M. Gastric bezoar: retrieve it, leave it, or disbelieve it?. Journal of Pediatric Gastroenterology and Nutrition, 2021, 72(2), e31-e36. https://doi.org/10.1097/MPG.0000000000002923.*

129. *Beyene, B.; Mohammadsani, B.; Abdlhadi, M.; Getachew, K.; Ahmedin, H. Khat induced bezoar; a rare cause of small bowel obstruction:a case report. Int J Surg Case Rep. 2022, 91, 106811. https://doi.org/10.1016/j.ijscr.2022.106811.*

130. *Mahir, S.; Salih, A.M.; Ahmed, O.F.; Kakamad, F.H.; Salih, R.Q.; Mohammed, S.H.; Usf, D.C.; Hassan, H.A.; Abdulla, F. Giant phytobezoar; an unusual cause of gastric outlet obstruction: A case report with literature review. Int J Surg Case Rep. 2020, 67, 154-156. https://doi.org/10.1016/j.ijscr.2020.02.012.*

131. *Doughty, M.; Alookaran, J.; Rhoads, M.; Dahm, P.H. Restrictive Lung Pattern: A Rare Complication of a Phytobezoar. JPGN Rep. 2021, 2(2):e056. https://doi.org/10.1097/PG9.0000000000000056.*

132. *Rahul, S.K.; Keshri, R.; Chaubey, D. Ileal bezoar secondary to ileal duplication - a case report. Trop Doct. 2025, 55(2), 179-181. https://doi.org/10.1177/00494755251319146.*
133. *Huang, Y.; Meng, J.; Zhang, S. Small intestinal obstruction caused by the gastric bezoars in elderly diabetic patients: A case report. Int J Surg Case Rep. 2025, 127, 110761. https://doi.org/10.1016/j.ijscr.2024.110761.*
134. *Lim, C.H.; Lim, C.J.; Yao, C.T.; Chang, C.C. Novel approach to managing two enormous bezoars with successive snare-tip electrocautery: A case report. World J Gastrointest Endosc. 2025, 17(1), 102185. https://doi.org/10.4253/wjge.v17.i1.102185.*
135. *Alouani H, Lâamiri G, Kallel Y, Arfaoui C, Bouassida M, Touinsi H. Gastrointestinal bezoars: two distinct cases illustrating diagnostic and surgical challenges. Int J Surg Case Rep. 2026 Feb 18;138(3):940-943. doi: 10.1097/RC9.0000000000000246. PMID: 41837025; PMCID: PMC12981764.*
136. *Angulo, S. D., Giraldo, Á. M., Sáenz Pinto, A., Vazquez, K. O., Mejía, S., & Rodriguez, M. S. (2026). Beyond usual suspects: phytobezoar in pediatrics a diagnostic challenge case report. International journal of surgery case reports, 138(4), 1322–1326. https://doi.org/10.1097/RC9.0000000000000198*
137. *Rojo, S. A. L., & Aguilera, J. F. C. (2026). Biliary phytobezoar associated with severe cholangitis. Cirujano General, 48(1), 40-46. doi:10.35366/123069.*
138. *Hegedűs, F., Lantos, T., & Sejben, A. (2026). Unexpected Findings on Histology: Plant Seeds Inducing and Mimicking Gastrointestinal Diseases. Diagnostics (Basel, Switzerland), 16(6), 826. https://doi.org/10.3390/diagnostics16060826*
139. *Hu, Z. J., Wang, Y., Ma, C. M., Zhou, H. H., & Jing, D. H. (2026). Case Report: Successful endoscopic treatment of gastric outlet obstruction due to duodenal persimmon phytobezoar. Frontiers in medicine, 13, 1732300. https://doi.org/10.3389/fmed.2026.1732300*
140. *Nawab, S., Shah, M. A., Nambiar, R., Longi, A. A., & Alexander, G. (2026). Phytobezoar-Induced Jejunal Obstruction in an Elderly Male With Multiple Comorbidities. Cureus, 18(3), e105285. https://doi.org/10.7759/cureus.105285*
141. *Ooya, Y., & Kaji, M. (2026). A case of small bowel obstruction caused by konjac. Journal of surgical case reports, 2026(4), rjag277. https://doi.org/10.1093/jscr/rjag277*
142. *Pandit, A., Koirala, M., Panta, A., & Kandel, N. (2026). Phytobezoar-An Unusual Cause of Small Bowel Obstruction in Pediatric Age Group: A Case Report and Literature Review. Clinical case reports, 14(1), e71809. https://doi.org/10.1002/ccr3.71809*
143. *Sanner Dixon, K., Nawabi, N., Hounschell, C., Cummiskey, B., Hass, S., McCoy, C., & Eaton, S. (2026). Small Bowel Obstruction Caused by a Bezoar in a Duodenal Diverticulum. Cureus, 18(1), e101549. https://doi.org/10.7759/cureus.101549*
144. *Valencia Valverde, S. A., Moyón Constante, F. X., & Loaiza Romero, S. P. (2026). Gastric coconut bezoar following Roux-en-Y gastric bypass. Journal of surgical case reports, 2026(5), rjag357. https://doi.org/10.1093/jscr/rjag357*
145. *Wu, X., Zhang, W., & Xia, M. (2026). Phytobezoar-induced intestinal obstruction relieved via enteroscopy. Revista espanola de enfermedades digestivas, 118(4), 234–235. https://doi.org/10.17235/reed.2024.10976/2024*
146. *Sani, A. V., Rashed, M. M., Atqiaee, K., Jafari, S. A., & Alamdaran, S. A. (2026). A Small Bowel Phytobezoar in a Child with Down Syndrome: Partial Response to Ultrasound-Guided Coca-Cola Injection. jpp, 5629.*
147. *Sabatella, L., Gascón Jiménez, J. J., Blanco, N., Rotellar, F., & Valentí, V. (2026). Disolución exitosa de un bezoar gigante tras tratamiento con complejo enzimático. Anales del sistema sanitario de Navarra, 49(1), e1154. https://doi.org/10.23938/ASSN.1154*
148. *Durán, R. B., & Rodrígez, J. R. L. (2026). Fitobezoar como causa inusual de obstrucción intestinal mecánica alta. Revista Cubana de Medicina Militar, 55(1), e026076870-e026076870.*
149. *Sotiropoulou, A., Martin, D., Vionnet, J., & Kefleyesus, A. Unusual Cause of Small Bowel Obstruction in a Liver Transplant Recipient: A Case Report and Focused Literature Review. Frontiers in Surgery, 13, 1824834.*
150. *Luyt, K. S., & Sukhera, F. (2026). Phytobezoar-induced small bowel obstruction in a child with Meckel's diverticulum. The American Journal of the Medical Sciences, 371, S467. https://doi.org/10.1016/j.amjms.2025.12.746.*

**S2. Appendix**

*151.* **PRISMA 2020 Checklist**

| Section and Topic | Item # | Checklist item | Reported on page |
|---|---|---|---|
| **TITLE** | | | |
| Title | 1 | Identify Systematic Review | 1 |
| **ABSTRACT** | | | |
| Abstract | 2 | A structured summary | 2 |
| **INTRODUCTION** | | | |
| Rationale | 3 | Describe the rationale for the review in the context of existing knowledge. | 3 |
| Objectives | 4 | Provide an explicit statement of the objective(s) or question(s) the review addresses. | 4,5 |
| **METHODS** | | | |
| Eligibility criteria | 5 | Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses. | 5,6 |
| Information sources | 6 | Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted. | 5 |
| Search strategy | 7 | Present the full search strategies for all databases, registers and websites, including any filters and limits used. | 5 |
| Selection process | 8 | Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process. | 6 |
| Data collection process | 9 | Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process. | 6 |
| Data items | 10a | List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect. | 5 |
| | 10b | List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information. | 5 |
| Study risk of bias assessment | 11 | Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process. | NA |
| Effect measures | 12 | Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results. | NA |

| Section and Topic | Item # | Checklist item | Reported on page |
|---|---|---|---|
| Synthesis methods | 13a | Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)). | 5,6 |
| | 13b | Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions. | 5,6 |
| | 13c | Describe any methods used to tabulate or visually display results of individual studies and syntheses. | 5,6 |
| | 13d | Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used. | 5,6 |
| | 13e | Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression). | NA |
| | 13f | Describe any sensitivity analyses conducted to assess robustness of the synthesized results. | NA |
| Reporting bias assessment | 14 | Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases). | NA |
| Certainty assessment | 15 | Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome. | 6 |
| **RESULTS** | | | |
| Study selection | 16a | Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram. | 7 |
| | 16b | Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded. | 7 |
| Study characteristics | 17 | Cite each included study and present its characteristics. | S1 appendix |
| Risk of bias in studies | 18 | Present assessments of risk of bias for each included study. | NA |
| Results of individual studies | 19 | For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/ credible interval), ideally using structured tables or plots. | S1 appendix |
| Results of syntheses | 20a | For each synthesis, briefly summarise the characteristics | S1 appendix |
| | 20b | Present results of all statistical syntheses conducted. | 7-9 |
| | 20c | Present results of all investigations of possible causes of heterogeneity among study results. | NA |
| | 20d | Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results. | NA |
| Reporting biases | 21 | Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed. | NA |
| Certainty of evidence | 22 | Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed. | 6 |
| **DISCUSSION** | | | |
| Discussion | 23a | Provide a general interpretation of the results in the context of other evidence. | 12-13 |
| | 23b | Discuss any limitations of the evidence included in the review. | 13 |
| | 23c | Discuss any limitations of the review processes used. | 13 |
| | 23d | Discuss implications of the results for practice, policy, and future research. | 12-13 |

| Section and Topic | Item # | Checklist item | Reported on page |
|---|---|---|---|
| **OTHER INFORMATION** | | | |
| Registration and protocol | 24a | Provide registration information for the review, including register name and registration number, or state that the review was not registered. | - |
| | 24b | Indicate where the review protocol can be accessed, or state that a protocol was not prepared. | NA |
| | 24c | Describe and explain any amendments to information provided at registration or in the protocol. | NA |
| Support | 25 | Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review. | 13 |
| Competing interests | 26 | Declare any competing interests of review authors. | 13 |
| Availability of data, code and other materials | 27 | Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review. | S1 appendix |

***NA: Not applicable**